



# **Shree Santkrupa College of Pharmacy, Ghogaon**

## **Criterion 3**

### **Research, Innovations and Extension**

#### **3.3**

##### **Research Publications and Awards**

---

##### **3.3.1**

**Number of research papers published per teacher in the Journals notified on UGC care list during the last five years**



**3.3.1 Number of research papers published per teacher in the Journals notified on UGC CARE list during the last five years**

Title of paper	Name of the author/s	Department of the teacher	Name of journal	Calendar Year of publication	ISSN number	Link to the recognition in UGC enlistment of the Journal /Digital Object Identifier (doi) number		
						Link to website of the Journal	Link to article / paper / abstract of the article	Is it listed in UGC Care list
Drug Delivery and Therapeutic Approaches to Prostate Cancer	<b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2018	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/drug-delivery-and-therapeutic-approaches-to-prostate-cancer/">https://www.ijndd.in/drug-delivery-and-therapeutic-approaches-to-prostate-cancer/</a>	Yes
Formulation and Standardization of Asava of Syzygium Cumini	<b>Patil S. V., Aralelimath V. R., Mahajan V. A., Inamdar N. R. and Shinde S. S.</b>	Pharmaceutics Pharmacology	Indian Drugs	2018	0019-462X	<a href="https://www.indiandrugsonline.org/">https://www.indiandrugsonline.org/</a>	<a href="https://www.indiandrugsonline.org/issues/article-details?id=ODM4">https://www.indiandrugsonline.org/issues/article-details?id=ODM4</a>	Yes
Formulation and Evaluation of Fast Dissolving Oral Films of Domperidone	<b>Jameel Ahmed S Mulla, Utkarsh A Chopade, Suraj B Kumbhar, Pallavi S Marathe, Priyanka V Ware</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2018	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/formulation-and-evaluation-of-fast-dissolving-oral-films-of-domperidone/">https://www.ijndd.in/formulation-and-evaluation-of-fast-dissolving-oral-films-of-domperidone/</a>	Yes
Effect of Polyethylene Glycol on	<b>Aralelimath V. R., Patil</b>	Pharmacology	Current Pharma Research	2018	2230-7842	<a href="https://jcpr.manjournals.com/">https://jcpr.manjournals.com/</a>	<a href="https://jcpr.manjournals.com/effect-of-">https://jcpr.manjournals.com/effect-of-</a>	Yes

Physiochemical Properties of Spherical Agglomerates of Antidiabetic Drug	<b>S. V., Patrakar R. G.</b>	Pharmaceutics Pharmacognosy					<a href="https://www.ijndd.in/development-and-characterization-of-rapidly-disintegrating-tablets-of-amlodipine-besylate/">polyethylene-glycol-on-physiochemical-properties-of-spherical-agglomerates-of-antidiabetic-drug/</a>	
Development and Characterization of Rapidly Disintegrating Tablets of Amlodipine Besylate	Shivani D Jadhav, <b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2018	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/development-and-characterization-of-rapidly-disintegrating-tablets-of-amlodipine-besylate/">https://www.ijndd.in/development-and-characterization-of-rapidly-disintegrating-tablets-of-amlodipine-besylate/</a>	Yes
Formulation and Evaluation of Chitosan Based Transdermal Patches of Lornoxicam for Prolonged Drug Release and to Study the Effect of Permeation Enhancer	Adhikrao Vyanktrao Yadav, <b>Mukund Namdeo Urade</b>	Pharmacology	Indian Journal of Pharmaceutical Education and Research	2019	0019-5464	<a href="https://www.ijper.org/">https://www.ijper.org/</a>	<a href="https://www.ijper.org/article/909">https://www.ijper.org/article/909</a>	Yes
Formulation and In Vitro, In Vivo Evaluation of Proniosomal Gel of Neomycin Sulphate	Amol Shete, <b>Priyanka Thorat,</b> Rajendra Doijad, Sachin Sajane	Pharmaceutics	International Journal of Applied Pharmaceutics	2019	0975-7058	<a href="https://journals.innovareacademics.in/index.php/ijap/index">https://journals.innovareacademics.in/index.php/ijap/index</a>	<a href="https://journals.innovareacademics.in/index.php/ijap/article/view/30614">https://journals.innovareacademics.in/index.php/ijap/article/view/30614</a>	Yes
Optimization of Thiazolidone Scaffolds Using Pocket Modeling for	Shivratna V. Khare, Sujata P. Choudhari, Siddharth P. Phalle, Santosh	Pharmaceutics	Turkish Journal of Pharmaceutical Sciences	2019	2148-6247	<a href="https://www.turkjps.org/">https://www.turkjps.org/</a>	<a href="https://www.turkjps.org/archives/archive-detail/article-preview/optimization-of-thiazolidone-scaffolds-">https://www.turkjps.org/archives/archive-detail/article-preview/optimization-of-thiazolidone-scaffolds-</a>	Yes

Development of Potential Secretary System Inhibitors of <i>Mycobacterium tuberculosis</i>	S. Kumbhar 1, Prafulla B. Choudhari, Sambhaji R. Masal, Aakash K. Patil, Rakesh P. Dhavale, Durgacharan A. Bhagwat, <b>Atul M. Kadam</b>						<a href="#">using-pocke/26306</a>	
Analytical method development and validation of Valacyclovir to estimate from its formulation	<b>P. D. Lade, S. V. Patil, S. S. Kadam, O. B. Tipugade, P. G. Nakhare</b>	Pharmaceutical Chemistry  Pharmaceutics	Current Pharma Research	2019	2230-7842	<a href="https://jcpr.manjournals.com/">https://jcpr.manjournals.com/</a>	<a href="https://jcpr.manjournals.com/analytical-method-development-and-validation-of-valacyclovir-to-estimate-from-its-formulation/">https://jcpr.manjournals.com/analytical-method-development-and-validation-of-valacyclovir-to-estimate-from-its-formulation/</a>	Yes
Design, Development and Characterization of Fast Dissolving Oral Film of Clonazepam	Govind G Nikam, <b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2020	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/design-development-and-characterization-of-fast-dissolving-oral-film-of-clonazepam/">https://www.ijndd.in/design-development-and-characterization-of-fast-dissolving-oral-film-of-clonazepam/</a>	Yes
A Novel Corticosteroid Cubosomes – For Ocular Drug Delivery	Snehal Shashaikant Chakorkar, <b>Jameel Ahmed S. Mulla</b>	Pharmaceutics	Indo American Journal of Pharmaceutical Research	2020	2231-6876	<a href="https://www.iajpr.com/">https://www.iajpr.com/</a>	<a href="https://zenodo.org/records/3923714">https://zenodo.org/records/3923714</a>	Yes
Formulation and Evaluation of Teneligliptin-Loaded	<b>Jameel Ahmed S Mulla, Vijayana and R Aralelim</b>	Pharmaceutics  Pharmacology	Indian Journal of Novel Drug Delivery	2020	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/formulation-and-evaluation-of-teneligliptin-loaded-">https://www.ijndd.in/formulation-and-evaluation-of-teneligliptin-loaded-</a>	Yes

Mucoadhesive Microspheres	<b>ath,</b> Omkar Tipugade, Shreyasi S Shinde, Nisha G Tetgure, Ayesha A Mulla, Dinesh D Gavali						<a href="#">mucoadhesive-microspheres/</a>	
Characterization, antioxidant, antimicrobial and cytotoxic activities of green synthesized silver and iron nanoparticles using alcoholic <i>Blumea ciliaris</i> D C plant extract	Chavan, R. R., Bhinge, S. D., Bhutkar, M. A., Randive, D. S., Wadkar, G. H., Todkar, S. S., & <b>Urade, M. N.</b>	Pharmacology	Materials Today Communications	2020	2352-4928	<a href="https://www.sciencedirect.com/journal/materials-today-communications">https://www.sciencedirect.com/journal/materials-today-communications</a>	<a href="https://www.sciencedirect.com/science/article/abs/pii/S235249282032331X">https://www.sciencedirect.com/science/article/abs/pii/S235249282032331X</a>	Yes
Microemulsion Based Hydrogel Formulation for Topical Drug Delivery - A Concise Review	<b>Jameel Ahmed S Mulla</b> and Biradev S Karande	Pharmaceutics	Indian Journal of Novel Drug	2021	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/microemulsion-based-hydrogel-formulation-for-topical-drug-delivery-a-concise-review/">https://www.ijndd.in/microemulsion-based-hydrogel-formulation-for-topical-drug-delivery-a-concise-review/</a>	Yes
Effect of polymers and process parameters in augmenting the compactability and dissolution behaviour of oxcarbazepine spherical	Sandip Honman, <b>Atul Kadam,</b> Sujata Choudhari, Raviraj Patil, Siddique Akber Ansari, Vinod Gaikwad	Pharmaceutics	Journal of Drug Delivery Science and Technology	2021	1773-2247	<a href="https://www.sciencedirect.com/journal/journal-of-drug-delivery-science-and-technology">https://www.sciencedirect.com/journal/journal-of-drug-delivery-science-and-technology</a>	<a href="https://www.sciencedirect.com/science/article/abs/pii/S1773224721002586">https://www.sciencedirect.com/science/article/abs/pii/S1773224721002586</a>	Yes

agglomerates								
Formulation and Characterization of Rapidly Dissolving Buccal Films of Montelukast Sodium	Ankita B Hogale, <b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug	2021	0975-5500	<a href="https://www.ijnidd.in/">https://www.ijnidd.in/</a>	<a href="https://www.ijnidd.in/formulation-and-characterization-of-rapidly-dissolving-buccal-films-of-montelukast-sodium/">https://www.ijnidd.in/formulation-and-characterization-of-rapidly-dissolving-buccal-films-of-montelukast-sodium/</a>	Yes
Ecopharmacovigilance : Perspectives, concepts, applications, and relationships in modern context	Prachi Khamkar, Debarshi KarMaha patra, <b>Atul Kadam</b>	Pharmaceutics	Journal of Pharmacovigilance and Drug Research	2021	2582-7235	<a href="https://www.jpadr.com/index.php/jpadr">https://www.jpadr.com/index.php/jpadr</a>	<a href="https://jpadr.com/index.php/jpadr/article/view/19">https://jpadr.com/index.php/jpadr/article/view/19</a>	Yes
Regulatory challenges and myths in pharmaceutical 3D printing	<b>Atul Kadam</b> , Prachi Khamkar	pharmaceutics	European Pharmaceutical Review	2021	1360-8606	<a href="https://www.europeanpharmaceuticalreview.com/">https://www.europeanpharmaceuticalreview.com/</a>	<a href="https://www.europeanpharmaceuticalreview.com/article/143008/regulatory-challenges-and-myths-in-pharmaceutical-3d-printing/">https://www.europeanpharmaceuticalreview.com/article/143008/regulatory-challenges-and-myths-in-pharmaceutical-3d-printing/</a>	Yes
Formulation and Process Validation of Clarithromycin Loaded Immediate Release Tablets	Ketaki S. Shinde, Dr. R. C. Doijad, <b>Dr. J. S. Mulla</b> and Sachin S. Mali	Pharmaceutics	World Journal of Pharmacy and Pharmaceutical Sciences	2022	2278-4357	<a href="https://www.wjpps.com/wjpps_controller/index">https://www.wjpps.com/wjpps_controller/index</a>	<a href="https://www.wjpps.com/Wjpps_controller/abstract_id/16195">https://www.wjpps.com/Wjpps_controller/abstract_id/16195</a>	Yes
Basic Aspects of Pharmaceutical Process Validation of Solid Dosage Forms: Quality Assurance	Ketaki S. Shinde, Dr. R. C. Doijad, <b>Dr. J. S. Mulla</b> and Sachin S. Mali	Pharmaceutics	European journal of pharmaceutical and medical research	2022	2394-3211	<a href="https://www.ejpmr.com/">https://www.ejpmr.com/</a>	<a href="https://www.ejpmr.com/home/abstract_id/9508">https://www.ejpmr.com/home/abstract_id/9508</a>	Yes

Point of View								
Floating Microspheres: A Novel Drug Delivery System	Omkar B Tipugade, <b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2022	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/floating-microspheres-a-novel-drug-delivery-system/">https://www.ijndd.in/floating-microspheres-a-novel-drug-delivery-system/</a>	Yes
<i>Periplaneta americana</i> L. a potential source of traditional medicine: chemometric analysis, <i>in vitro</i> and <i>in silico</i> study	Suchismeta Behera, Amulyarantna Behera, Suman Kumar Mekap, Chinmaya Chidananda Behera, <b>Atul Kadam &amp; Prafulla K. Mohanty</b>	Pharmaceutics	Journal of Biomolecular Structure and Dynamics	2022	1538-0254	<a href="https://www.tandfonline.com/journals/tbsd20">https://www.tandfonline.com/journals/tbsd20</a>	<a href="https://www.tandfonline.com/doi/abs/10.1080/07391102.2021.1938681">https://www.tandfonline.com/doi/abs/10.1080/07391102.2021.1938681</a>	Yes
Non-Ionic Surfactant Vesicle (Niosome): A Novel Drug Delivery System	Sandhya S Shewale, <b>Jameel Ahmed S Mulla</b>	Pharmaceutics	Indian Journal of Novel Drug Delivery	2022	0975-5500	<a href="https://www.ijndd.in/">https://www.ijndd.in/</a>	<a href="https://www.ijndd.in/non-ionic-surfactant-vesicle-niosome-a-novel-drug-delivery-system/">https://www.ijndd.in/non-ionic-surfactant-vesicle-niosome-a-novel-drug-delivery-system/</a>	Yes
Nanosuspension: A Novel Technology for Drug Delivery	Vidya Ashok Kheradkar, <b>Jameel Ahmed S. Mulla</b>	Pharmaceutics	Asian Journal of Research in Pharmaceutical Sciences	2023	2231–5640	<a href="https://ajpsonline.com/">https://ajpsonline.com/</a>	<a href="https://ajpsonline.com/AbstractView.aspx?PID=2023-13-2-6">https://ajpsonline.com/AbstractView.aspx?PID=2023-13-2-6</a>	Yes
Evaluation of solid-lipid nanoparticles formulation of	Debarati Maiti, Mohammed Naseeruddin Inamdar ,	Pharmaceutics	Saudi Pharmaceutical Journal	2023	1319-0164	<a href="https://www.sciencedirect.com/science/article/pii/S1319016423000890">Saudi Pharmaceutical Journal   ScienceDirect .com by Elsevier</a>	<a href="https://www.sciencedirect.com/science/article/pii/S1319016423000890">https://www.sciencedirect.com/science/article/pii/S1319016423000890</a>	Yes

methotrexate for anti-psoriatic activity	Sarasija Suresh, Syed Mohammed Basheeruddin Asdaq, <b>Jameel Ahmed S. Mulla</b> et al.							
Cubosomal Drug Delivery System: Promising Approach for Ocular Drug Delivery of Corticosteroid	Snehal Chakorkar, <b>Jameel Ahmed S. Mulla</b>	Pharmaceutics	European Chemical Bulletin	2023	2063-5346	<a href="https://www.eurchembull.com/">https://www.eurchembull.com/</a>	<a href="https://www.eurchembull.com/issue-content/cubosomal-drug-delivery-system-promising-approach-for-ocular-drug-delivery-of-corticosteroid-5440">https://www.eurchembull.com/issue-content/cubosomal-drug-delivery-system-promising-approach-for-ocular-drug-delivery-of-corticosteroid-5440</a>	Yes
Phytochemical and Pharmacological Review on <i>Garcinia talbotii</i> : A Latent Medicinal Plant	<b>Jameel Ahmed S. Mulla</b>	Pharmaceutics	Indian Journal of Natural Sciences	2023	0976 – 0997	<a href="https://tnsroindia.org.in/journals.html">https://tnsroindia.org.in/journals.html</a>	<a href="https://tnsroindia.org.in/JOURNAL/issue80/ISSUE%2080%20-%20OCTOBER%202023%20-%20FULL%20TEXT%2002.pdf">https://tnsroindia.org.in/JOURNAL/issue80/ISSUE%2080%20-%20OCTOBER%202023%20-%20FULL%20TEXT%2002.pdf</a>	Yes
Formulation and Evaluation of Poly Herbal Chewable Tablets from Ayurvedic Ingredients having Cough Relieving Activity	Vinayak Dasharath Gaikwad, <b>Priyanka A. Thorat</b>	Pharmaceutics	International Journal of Innovative Science and Research Technology	2023	2456-2165	<a href="https://ijisrt.com/">https://ijisrt.com/</a>	<a href="https://www.ijisrt.com/formulation-and-evaluation-of-poly-herbal-chewable-tablets-from-ayurvedic-ingredients-having-cough-relieving-activity">https://www.ijisrt.com/formulation-and-evaluation-of-poly-herbal-chewable-tablets-from-ayurvedic-ingredients-having-cough-relieving-activity</a>	Yes
Extraction,	<b>Pravinku</b>	Pharmaceutics	International	2023	0975	<a href="https://ijpqa.com">https://ijpqa.com</a>	<a href="https://impactf">https://impactf</a>	Yes



Phytochemical Screening and HPTLC Analysis of Fenugreek Gum for Analysis of Different Phytoconstituents	<b>Dr. D. Lade,</b> Neelam Singla	Journal of Pharmaceutical Chemistry	Journal of Pharmaceutical Quality Assurance		9506	<a href="#">http://www.ijpqa.com/</a>	<a href="http://www.ijpqa.com/PDF/IJPOA/14/IJPOA_Vol14_Issue4_Article42.pdf">http://www.ijpqa.com/PDF/IJPOA/14/IJPOA_Vol14_Issue4_Article42.pdf</a>	
---	--------------------------------------	-------------------------------------	---	--	------	---------------------------------------	---	--



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Review Article

**Drug Delivery and Therapeutic Approaches to Prostate Cancer**

JAMEEL AHMED S MULLA\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Tal- Karad, Dist- Satara, Maharashtra, India

**ARTICLE DETAILS**

*Article history:*

Received on 05 September 2018

Modified on 25 September 2018

Accepted on 27 September 2018

*Keywords:*

Prostate structure,  
Prostate cancer,  
Therapeutic approaches,  
Drug delivery approaches,  
Liposomes,  
Nanoparticles

**ABSTRACT**

According to the global cancer statistics, prostate cancer is the second most often diagnosed cancer in males worldwide and it ranks in fifth place among cancer-related deaths. Localized prostate cancer is usually treated by surgical removal of the prostate (radical prostatectomy) or by radiation therapy. Both treatment options are frequently associated with severe side effects. Drug delivery to prostate through conventional route is associated with pharmacokinetics based and side effects related problems. This review briefly highlighted about prostate structure and its cancer. Different therapeutic approaches such as radiotherapy, chemotherapy, immunotherapy, androgen deprivation therapy (ADT), focal therapies and systemic therapies; and drug delivery approaches such as liposomes, polymeric based nanoparticles and micelles, dendrimers, gold nanoparticles, carbon based nanoparticles and magnetic nanoparticles for prostate cancer are also outlined here.

© KESS All rights reserved

**INTRODUCTION  
Cancer**

Cancer is the second leading cause of death worldwide, and is accountable for a predictable 9.6 million deaths in 2018. Globally, about 1 in 6 deaths is due to cancer. About 70% of deaths from cancer occur in low- and middle-income countries. The most common cancers are; lung (2.09 million cases), breast (2.09 million cases), colorectal (1.80 million cases), prostate (1.28 million cases), skin cancer (non-melanoma) (1.04 million cases) and stomach (1.03 million cases). Cancer arises from the transformation of normal cells into tumour cells in a multistage process that normally progresses from a pre-cancerous lesion to a malignant tumour. These changes are the consequence of the interaction between a person's genetic factors and 3 categories of external agents, including (a) physical carcinogens, such as ultraviolet and ionizing radiation (b) chemical carcinogens, such as asbestos, components of tobacco smoke, aflatoxin (a food contaminant), and arsenic (a drinking water contaminant) (c) biological carcinogens, such as infections from certain viruses, bacteria, or parasites [1].

**Prostate**

The prostate is one of the male accessory glands that contribute to the male ejaculate by producing seminal fluid. In mice, elimination of this organ reduces fertility [2]. The understanding of development, maturation and anatomy of prostate has progressed gradually throughout the past century, but it has been moderately hindered by reliance on historical anatomical descriptions [3-5]. Description of prostatic lobes has been replaced by the now widely accepted concept that the human prostate is better described as consisting of zones found within one discrete organ [6].

The prostate is an exquisitely steroid hormone sensitive organ. Considering hormone action throughout the different life stages of prostate organogenesis, puberty, and during adulthood and aging, is critical for understanding prostate anatomy and the different diseases of this organ. Certainly, most of men experience some form of prostate pathology during their lifetime. The focus has been on investigating the long-term impact of disruption of normal prostate development on structural and functional abnormalities related to specific types of prostate disease that often are not recognized until much later in life [7].

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com



*(Handwritten signature)*  
**Principal**

**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Drug delivery and therapeutic approaches to prostate cancer

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 10

July-September 2018

Drug delivery and therapeutic approaches to prostate cancer

Jameel Ahmed S Mulla

Download PDF


**Abstract:**  
 According to the global cancer statistics, prostate cancer is the second most often diagnosed cancer in males worldwide and it ranks in fifth place among cancer-related deaths. Localized prostate cancer is usually treated by surgical removal of the prostate (radical prostatectomy) or by radiation therapy. Both treatment options are frequently associated with severe side effects. Drug delivery to prostate through conventional route is associated with pharmacokinetics based and side effects related problems. This review briefly highlighted about prostate structure and its cancer. Different therapeutic approaches such as radiotherapy, chemotherapy, immunotherapy, androgen deprivation therapy (ADT), local therapies, and systemic therapies, and drug

2 new notifications

Type here to search

25°C Sunny 12:20 03-10-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## FORMULATION AND STANDARDIZATION OF ASAVA OF SYZYGIIUM CUMINI

### ABSTRACT

Asavas and Arishta are alcoholic medicaments prepared by allowing the herbal juices or their decoctions to undergo fermentation with the addition of sugar. Standardization of ayurvedic formulation is essential in order to assess the quality of drugs. In the present study standardization of asava from *Syzygium cumini*, known to be effective in diabetics has been performed. Asava formulation was prepared by reported traditional method of ayurveda. Formulation has been standardized by modern scientific quality control procedure for the finished product. Standardization of asava was achieved by organoleptic study, physicochemical parameters such as pH, specific gravity, total solid content, acid value, alcohol content, refractive index, total reducing sugars and stability study. The results have revealed that the physicochemical parameters were within the limits and the values could be used to establish and formulate procedures for standardization and quality controlling of these ayurvedic formulations.

**Keywords:** standardization, asava, arishta, *syzygium cumini*, diabetics

The word Ayurveda is composed of two parts ayur and veda, Ayur means life and veda means knowledge. Ayurveda has long and strong heritage of use of poly herbal drugs and formulations to treat various diseases<sup>1</sup>. Asava and Arishta are alcoholic medicaments prepared by allowing the herbal juices or their decoctions to undergo fermentation with the addition of sugars<sup>2</sup>. Formulation of Asava and Arishta has done by soaking the drugs, either in coarse powder form or in the form of decoction, in a solution of sugar or jaggery, for a specified period of time, during which it undergoes a process of fermentation<sup>3-4</sup>. Asava have been used as medicines far over 3000 years to treat various disorders & are also taken as appetizers and stimulants. It is liquid preparations containing self generated alcohol, thus contain water soluble as well as alcohol soluble substances of the drugs. Due to their medicinal value, sweet taste, and easy availability people are prone to consume higher doses of these drugs for longer periods. As per Samhita, Asava is Madya which is prepared with apakwa Aushada. The compound which is prepared by 'asutaprakriya' is called as Asava<sup>5,6</sup>. Asava is a type of Madya, which contains decoction of different drugs, guda, and dhataki, described by Sushruta. Asava is defined as that which is prepared from ikshurasa, described by Sushruta. Vagbhata has also defined Asava as Madya prepared using fresh tubers, roots, fruits etc. and Madya with medicinal properties is called as Asava<sup>7-9</sup>. But the systematic quality control parameters of such problem is main reason for reproducibility along with control over batch to batch uniformity<sup>8, 10</sup>. The aim of present study was formulation and standardization of the Asava an Ayurvedic formulation to establish typical quality control parameters of such formulations.

All powdered ingredients were collected from Unique Chemicals, Kolhapur, Maharashtra, India. All Chemicals used were of analytical grade.

Approximately 250 mL of water taken in to the cleaned fermentation flask (glass; 350 mL), with hot water and Sugar/Jaggery/Wheat flour, was dissolved in to it. This mixture was boiled for half an hour, cooled to room temperature to have a final mixture. All ingredients as given in Table I were weighed and passed through sieve no 44 and ingredients from 1 to 14 were added to the final mixture. Dhataki pusha (*Woodfordia fruticosa*) was then added to the mixture. The vessel was closed with a clean lid followed by wrapping around the lid with clay smeared cloth. The vessel was kept at dark place for 40 days.

After the stipulated period, the vessel was withdrawn. Preparation showed dark brownish fluid, pleasant smell and alcoholic taste. The fluid was filtered through muslin cloth, prepared asava was kept in a container.

Organoleptic characteristics odour, taste, colour and clarity of prepared asava formulation was determined.

Digital pH meter was used to check the pH of formulations. Before the experiment the machine was calibrated by using standard buffer solution of pH 4.0, 7.0 and 9.2.


The density and specific gravity were determined by using density bottle method<sup>11</sup>.

The viscosity was determined by Ostwald viscometer.

### Total solid content determination

20 mL of the Asava formulation was taken in evaporating dish and evaporated to dryness on a water bath and then



  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Issue's Article Details


indiandrugsonline.org/issue/article-details/Tar-ODM4

INDIAN DRUGS

Home | About us | Advertise with us | Editorial Board | Subscribe | Contact

LOG IN REGISTER

Current Issue | Past Issues | Best Paper Awards | Articles Accepted | Instructions To Authors | **SUBMIT ARTICLE**

Follow Us  **Article Details**

**FORMULATION AND STANDARDIZATION OF ASAVA OF SYZYGIUM CUMINI**

Patil S. V.<sup>1</sup>, Anandkarli H. N.<sup>2</sup>, Mahajan V. A.<sup>3</sup>, Bhande H. N.<sup>4</sup> and Shinde S. S.<sup>5</sup>

<sup>1</sup> Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Satara - 415 111, Maharashtra, India  
<sup>2</sup> Anandkarli Naina College of Pharmacy, Peth Ghogaon, Tal- Hama Nagar, Dist- Kolhapur - 416 112, Maharashtra, India

\* For correspondence: E-mail - santkrupa11@gmail.com  
<https://doi.org/10.33794/ID.55.28.11079>

**ABSTRACT**

Asavas and Arishtas are alcoholic preparations are prepared by allowing the herbal parts in their decoctions to undergo fermentation with the addition of sugar. Standardization of ayurvedic formulation is essential in order to assess the quality of drugs. In the present study standardization of asava from Syzygium cumini, known to be effective in diabetes, has been performed. Asava formulation was prepared by reported traditional method of ayurveda. Formulation has been standardized by modern scientific quality control procedure for the finished products. Standardization of asava was achieved by organoleptic study, physicochemical parameters such as pH, specific gravity, total solid content, acid value, alcohol content, refractive index, total reducing sugars and stability study. Results reveal that the physicochemical parameters were within the limits and the values could be used to establish and formulate procedures for standardization and quality controlling of these ayurvedic formulations.

Nov 2019 | Volume No. 55 | Issue No. 08 | Page No. 83-85

**DOWNLOAD ARTICLE**

Recent Issue

November 2023  
Vol. 55, Issue 11

October 2023  
Vol. 55, Issue 10

September 2023  
Vol. 55, Issue 9




August 2023  
Vol. 55, Issue 8

View All


Current Issue

INDIAN DRUGS

December 2023

Follow Us   



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara





Research Article

**Formulation and Evaluation of Fast Dissolving Oral Films of Domperidone**

JAMEEL AHMED S MULLA\*, UTKARSH A CHOPADE, SURAJ B KUMBHAR, PALLAVI S MARATHE,  
PRIYANKA V WARE

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Tal:Karad, Dist: Satara, Maharashtra, India

**ARTICLE DETAILS**

*Article history:*  
Received on 15 April 2018  
Modified on 10 June 2018  
Accepted on 17 June 2018

*Keywords:*  
Fast Dissolving Oral Film,  
Domperidone,  
FTIR,  
DSC,  
Mouth Dissolving Time

**ABSTRACT**

The oral route remains the most preferred for the general population. This system will provide maximum therapeutic efficacy, increased bioavailability and maximum stability by reducing the frequency of dosage. The present work aims to formulate and evaluate fast dissolving oral films of domperidone using HPMC K100 and HPMC E15 as film forming agents by solvent casting method. The prepared films were characterized for weight variation, thickness, tensile strength, folding endurance, surface pH, *in vitro* disintegration time and mouth dissolving time. Polymer ratio and concentration of superdisintegrants played important role in disintegration and mouth dissolving time. *In vitro* drug release profile exhibited that formulation with higher concentration of superdisintegrants dissolves the oral films faster than the others.

© KESS All rights reserved

**INTRODUCTION**

Fast dissolving oral films provide the opportunity to administer medicines and avoid first-pass metabolism [1]. Fast dissolving oral films may also be used in children [2-3], patients with dysphasia [4], and elderly patients [5]. Although certain products such as paracetamol are available as oral suspension, these contain additives and sugar, which may not be advisable for children [6]. In addition, administering oral liquid formulations to children is challenging by using syringes [7]. These concerns are triggers for the development of more number of fast dissolving oral films formulations. A fast dissolving oral film has been successfully used to deliver medicines to patients having difficulty in swallowing, those with oral pain due to mucositis or after oral surgery, or those with nausea [8].

The oral route is one of the most preferred routes of drug administration as it is more convenient, cost effective, and ease of administration lead to high level of patient compliance. Recently, fast dissolving films are gaining interest as an alternative of fast dissolving tablets. The films are designed to dissolve upon contact with a wet surface, such as the tongue, within a few seconds, meaning the

consumer can take the product without need for additional liquid. It gives quick absorption and instant bioavailability of drugs due to high blood flow and permeability of oral mucosa is 4-1000 times greater than that of skin. This convenience provides both a marketing advantage and increased patient compliance. As the drug is directly absorbed into systemic circulation, degradation in gastrointestinal tract and first pass effect can be avoided. These points make this formulation most popular and acceptable among paediatric and geriatric patients and patients with fear of choking. Over-the-counter films for pain management and motion sickness are commercialized in the US markets.

Fast dissolving films are useful in patients such as paediatric, geriatrics, bedridden, emetic patients, diarrhoea, sudden episode of allergic attacks, or coughing for those who have an active life style. It is also useful whether local action desired such as local anaesthetic for toothaches, oral ulcers, cold sores or teething.

Advantages of fast dissolving films include; Convenient dosing, No water needed, No risk of choking, Taste masking, Enhanced stability, Improved patient compliance, The drug enters the systemic circulation with reduced hepatic first pass effect, Site specific and local action, Availability of large surface area that leads to rapid disintegration and dissolution within oral

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

68



*Principal*

**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Formulation and evaluation of...

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 10

April-June 2018

Formulation and evaluation of fast dissolving oral films of domperidone


Jameel Ahmed S Mulla, Ulkarsh A Chopade, Suraj B Kumbhar, Pallavi S Marathe, Priyanka Y Ware

Download PDF

**Abstract:**  
 The oral route remains the most preferred for the general population. This system will provide maximum therapeutic efficacy, increased bioavailability and maximum stability by reducing the frequency of dosage. The present work aims to formulate and evaluate fast dissolving oral films of domperidone using HPMC K100 and HPMC E15 as film forming agents by solvent casting method. The prepared films were characterized for weight variation, thickness, tensile strength, folding endurance, surface pH, in vitro disintegration time and mouth dissolution time. Release rate and permeation of domperidone through polymeric membrane was also studied.

25°C Sunny 12:27 03-02-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara

Research Article

**Effect of Polyethylene Glycol on Physiochemical Properties of Spherical Agglomerates of Antidiabetic Drug.**

**Aralelimath V. R.\* Patil S. V., Patrakar R. G.**

Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Dist. Satara, Maharashtra. India.

Received 04 March 2018; received in revised form 27 March 2018; accepted 27 March 2018

\*Corresponding author E-mail address: vijayananda\_15@yahoo.com

**ABSTRACT**


Spherically agglomerated crystals of Pioglitazone hydrochloride (PGH) with improved flowability and compactibility were successfully prepared by emulsion solvent diffusion method. Plane agglomerates and agglomerates with polyethylene glycol 6000 (PEG) 1% and 2% were prepared using methanol, chloroform and water as good solvent, bridging liquid and poor solvent respectively. Particle size, flowability, compactibility and packability of agglomerates were preferably improved for direct tableting compared with raw PGH. These improved properties of spherically agglomerated crystals were due to their large and spherical shape and enhanced fragmentation during compaction which was well supported by increased tensile strength and less elastic recovery of its compact. X-ray powder diffraction and differential scanning calorimetry study were indicated polymorphic transition of PGH from form II to I during recrystallization but not associated with chemical transition indicated by Fourier transforms infrared spectra.

**KEYWORDS**

Spherical crystallization, polyethylene glycol 6000, pioglitazone hydrochloride, compatibility, pack ability.

2368



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Effect of Polyethylene Glycol on Physiochemical Properties of Spherical Agglomerates of Antidiabetic Drug

JCPR Journal of Current Pharma Research  
 (An Official Publication of Human Journals)  
 An International Peer Reviewed Journal For Pharmacy, Medical & Biological Science  
 DOI: 10.25166 CODEN: JCPRD6 NLM ID: 101744065  
 ISSN Print: 2250-7834  
 Online: 2750-7847  
 SJIF Impact Factor: 8.332

Home About Us Instructions to Author Publication Fee Manuscript Submission Archives Contact Us

Effect of Polyethylene Glycol on Physiochemical Properties of Spherical Agglomerates of Antidiabetic Drug

MARCH 18, 2018

DOWNLOAD ARTICLE HERE: 8, Anandlimath V. R., Patil S. V., Patrakar R. G. (803.39 kB)

Next Story: A Review Transdermal Microneedle

Previous Story: An Overview on Validation Technique

Search

SJIF IMPACT FACTOR 8.332

Scientific Journal  
**8.332**  
 Impact Factor

Type here to search 26°C Sunny 12:00 03-02-2024



*[Handwritten Signature]*  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## Research Article

**Development and Characterization of Rapidly Disintegrating Tablets of Amlodipine Besylate**

SHIVANI D JADHAV, JAMEEL AHMED S MULLA\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Tal- Karad, Dist- Satara, Maharashtra, India

**ARTICLE DETAILS***Article history:*

Received on 07 May 2018

Modified on 20 June 2018

Accepted on 24 June 2018

*Keywords:*Amlodipine Besylate,  
Rapidly Disintegrating Tablets,  
Superdisintegrants,  
Disintegration Time,  
In Vitro Drug Release**ABSTRACT**

Hypertension is one of the most important causes of premature death worldwide and the problem is growing; in 2025, an estimated 1.56 billion adults will be living with hypertension. Amlodipine besylate is a longacting calcium channel blocker used in the treatment of chronic stable angina, vasospastic angina and hypertension. Recent developments in the technology have prompted scientists to develop Rapidly Disintegrating Tablets (RDTs) with improved patient compliance and convenience. The present study aims to prepare and evaluate amlodipine besylate Rapidly Disintegrating Tablets using superdisintegrants by direct compression method. Amlodipine loaded-RDTs were characterized for weight variation, hardness, thickness, friability, wetting time and disintegration time. In vitro drug release study was performed using United States Pharmacopoeia (USP) II dissolution testing apparatus II (paddle method). The results revealed that Amlodipine loaded-RDTs were successfully formulated with good mouth feel, faster disintegration and better drug release.

© KESS All rights reserved

**INTRODUCTION**

Hypertension, also known as high or raised blood pressure, is a condition in which the blood vessels have persistently raised pressure. Blood is carried from the heart to all parts of the body in the vessels. Each time the heart beats, it pumps blood into the vessels. Blood pressure is created by the force of blood pushing against the walls of blood vessels (arteries) as it is pumped by the heart. The higher the pressure the harder the heart has to pump [1].

Globally, nearly one billion people have high blood pressure (hypertension); of these, two-thirds are in developing countries. Hypertension is one of the most important causes of premature death worldwide and the problem is growing; in 2025, an estimated 1.56 billion adults will be living with hypertension. Hypertension kills nearly 8 million people every year, worldwide and nearly 1.5 million people each year in the South-East Asia (SEA) Region. Approximately one-third of the adult population in the SEA Region has high blood pressure [2].

Amlodipine besylate, chemically described as 3-Ethyl-1-methyl (±)-2-[[2-aminoethoxy] methyl]-4-(2-chlorophenyl)-1, 4-dihydro -6-methyl-3, 5-pyridinedicarboxylate monobenzenesulphonate [3-4], is a long-acting dihydropyridine calcium channel blocker commonly used in the treatment of hypertension. Amlodipine selectively inhibits transmembrane influx of calcium ions into cardiac and vascular smooth muscle, leading to decreased vascular tone, reduced systemic vascular resistance, diminished after load and coronary vasodilation [5].


The term oral drug delivery, also known as peroral delivery, refers to taking a dosage form by mouth for local action or systemic absorption at any point along the gastrointestinal (GI) tract. Oral drug delivery is the most readily available and widely accepted route of delivery for medications. The most obvious challenge is that many drugs have a bad taste when placed in the mouth or exhibit foul odors, which must be resolved to be acceptable to patients. Apart from this, there are also many barriers and obstacles that inhibit drug absorption via the oral route. These include the acidic gastric environment, digestive enzymes, mucus layer diffusion and tight junctions between epithelial cells that prevent paracellular transport [6-7]. Many

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

84



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Development and characterization of rapidly disintegrating tablets of amlodipine besylate

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 10

April-June 2018

← Back to Issue

Development and characterization of rapidly disintegrating tablets of amlodipine besylate

Shivani D. Jadhav, Jameel Ahmed S. Mulla


Download PDF

**Abstract:**  
Hypertension is one of the most important causes of premature death worldwide and the problem is growing. In 2025, an estimated 1.56 billion adults will be living with hypertension. Amlodipine besylate is a long acting calcium channel blocker used in the treatment of chronic stable angina, vasospastic angina and hypertension. Recent developments in the technology have prompted scientists to develop Rapidly Disintegrating Tablets (RDTs) with improved patient compliance and convenience. The present study aims to prepare and evaluate amlodipine besylate Rapidly Disintegrating Tablets using superdisintegrants by direct compression method.

Type here to search

25°C Sunny 12:31 03-02-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



# Formulation and Evaluation of Chitosan Based Transdermal Patches of Lornoxicam for Prolonged Drug Release and to Study the Effect of Permeation Enhancer

Adhikrao Vyankatrao Yadav<sup>1</sup>, Mukund Namdeo Urade<sup>2\*</sup>

<sup>1</sup>Gourishankar Institute of Pharmaceutical Education and Research, Limb, Satara- 415002, Maharashtra, INDIA.

<sup>2</sup>Research Scholar, Government College of Pharmacy, Karad - 415110, Maharashtra, INDIA.

## ABSTRACT

**Objective:** The objectives of present investigation were to develop transdermal system for Lornoxicam using chitosan as rate controlling polymer and Tween 20 as permeation enhancer. Then evaluate the effect of Tween 20 on the physico-chemical properties of the patches and on drug permeation across the membrane. **Methods:** Transdermal patches of Lornoxicam were prepared by solvent casting method. The prepared patches were evaluated for physicochemical characteristics such as *in vitro* drug release, skin irritation studies. The interaction between drug and polymer were investigated by FTIR, DSC, XRPD methods. **Results:** The *in vitro* release studies revealed Formulation L4 containing higher concentration of Tween 20 showed 74.6% diffusion in 12 h and follows Korsmeyer-Peppas drug release kinetics. Respective formulation did not showed any sign of erythema or edema in skin irritation test. FTIR study reveals good compatibility between drug and polymer. **Conclusion:** The prepared transdermal drug delivery system of Lornoxicam using Chitosan had shown good promising results for sustained release matrix transdermal patch formulation.

**Key words:** Transdermal patches, Chitosan, Lornoxicam, Tween 20, *in-vitro* release studies.

## INTRODUCTION

Transdermal drug delivery system provides controlled drug release through skin to reach systemic circulation. Transdermal delivery also has leading edge over oral route by avoiding first pass metabolism of drugs. Transdermal delivery serves variety of advantages compared with the traditional oral route of drug administration. Transdermal drug delivery systems are non-invasive and inexpensive, self-administered, improve patient compliance and can provide release of drugs for long periods of time. The challenge remains for transdermal delivery is that only limited number of drugs is amenable to administration by this route.<sup>1</sup> Treatment of chronic diseases/disorders such as asthma, rheumatoid arthritis by transdermal route of drug administration

might prove to have several advantages over other routes of drug administration.<sup>2</sup>

Chitosan is N-deacetylated derivative of chitin, semi-rigid polysaccharide, which is composed of N-acetyl D-glucosamine and D-glucosamine. Chitosan is a promising biomaterial in view of its biodegradability; biocompatibility and nonantigenicity.<sup>3</sup> Chitosan has bio-adhesive or mucoadhesive property that readily binds to negatively charged surfaces such as mucosal membranes. Chitosan greatly enhances the transport of polar drugs across epithelial surfaces. Chitosan is insoluble in neutral and basic environments due to lack of positive charge. Chitosan has a positive charge under acidic conditions due to protonation of free amino groups which enhances the solubility.<sup>4</sup>

Submission Date: 31-03-2018;

Revision Date: 14-08-2018;

Accepted Date: 23-10-2018

DOI: 10.5530/ijper.53.1.12

Correspondence:

Mr. Mukund N. Urade,  
Government College of  
Pharmacy, Vidyanagar, Taluka  
Karad-41511, District Satara,  
Maharashtra, INDIA.  
Phone: +91 9890111816  
E-mail: mukundurade@  
gmail.com



www.ijper.org



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Formulation and Evaluation of ...

ijper.org/article/309

**IJPER** INDIAN JOURNAL OF PHARMACEUTICAL EDUCATION AND RESEARCH

Abstract and Indexed in: *Journal of Indian Medicines Education, Journal of Indian Pharmacy, IJPER, IJPER, IJPER*

*Quality Publications Since 1967*

An official publication of Association of Pharmaceutical Teachers of India (APTI) ISSN: 0019-5464

Home About APTI About Journal For Authors For Librarians Search Articles Contact Us

Home | Articles

Tue 11/27/2019 09:42 | anonymous

How Pharmacy Students Evaluate the Credibility of Scientific Information: A Qualitative Study

Challenges in Simultaneous Analysis of Hydrophilic and Lipophilic Drugs in Combined Dosage Forms

Published on January 2019  
 Indian Journal of Pharmaceutical Education and Research, 2019, 53(1):99-99  
 Original Article | doi:10.5530/ijper.53.1.12

**Formulation and Evaluation of Chitosan Based Transdermal Patches of Lornoxicam for Prolonged Drug Release and To Study the Effect of Permeation Enhancer**

**Authors and affiliation (s):**  
 Adhikar V, Nandkumar V, Mahesh Kumar C

1. Government Institute of Pharmaceutical Education and Research, Pimpri, Satara, Maharashtra, 421004

Browse Issues

In Press

Latest Issue

Peer Review

RSS Feeds

Impact Factor


IJPER is Official Publication of Association of Pharmaceutical Teachers of India. It is granted to Associate Professor in the IJPER Review of Journal Chapter Reports (online Web of Science Data).

Impact Factor: as reported in the 2018 Journal Citation Reports® (Clarivate Analytics, 2019): 3.8

Type here to search

27°C Sunny 12:37 25-10-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara

## FORMULATION AND *IN VITRO*, *IN VIVO* EVALUATION OF PRNOSOMAL GEL OF NEOMYCIN SULPHATE

AMOL SHETE<sup>1</sup>, PRIYANKA THORAT<sup>2</sup>, RAJENDRA DOJJAD<sup>3</sup>, SACHIN SAJANE<sup>3</sup>

<sup>1</sup>Department of Pharmaceutics, Krishna Institute of Pharmacy of Krishna Institute of Medical Sciences Deemed To Be University, Karad, 415110, India, <sup>2</sup>Research Group Department of Pharmaceutics and Quality Assurance, Shree Santkrupa College of Pharmacy, Ghogaon, Karad 415111, MS, India, <sup>3</sup>Department of pharmacology, Annasaheb Dange College of pharmacy, Ashta, Maharashtra, India  
Email: amol.shete@rediffmail.com

Received: 01 Nov 2018, Revised and Accepted: 30 Jan 2019

### ABSTRACT

**Objective:** The objectives of present investigation were to prepare and evaluate proniosomes of neomycin sulphate (NS) by coacervation phase separation method by using sorbitan monostearate (span 60) and lecithin as a surfactant to increase the penetration through the skin and study the effect of concentration of the same.

**Methods:** Proniosomes of neomycin sulphate (NS) were prepared by coacervation phase separation method by using span 60 and lecithin. The effect of concentration of span 60 and lecithin was studied by factorial design. The prepared proniosomes were converted to gel by using carbopol as a gelling agent. The prepared formulations were evaluated for entrapment efficiency, *in vitro* drug diffusion, *in vitro* antibacterial activity and *in vivo* skin irritation test etc.

**Results:** All Formulation showed the percentage entrapment efficiency in the range 38.31±0.05% to 77.96±0.06%, good homogeneity and gel was easily spreadable with minimal of shear. Optimized formulation showed enhanced rate of diffusion *in vitro*, increase in zone of inhibition against *staphylococcus aureus*, no skin irritation and showed good stability.

**Conclusion:** The results of present study indicates that proniosomal gel formulated by using combination of span 60, Lecithin, cholesterol can be used to enhance skin delivery of NS because of excellent permeation of drug. Developed proniosomal gel formulation was promising carrier for NS

**Keywords:** Proniosomes, Neomycin Sulphate, Sorbitane monostearate (span 60), Lecithin and Cholesterol

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)  
DOI: <http://dx.doi.org/10.22159/ijap.2019r11i2.30614>

### INTRODUCTION

Oral route of administration is most accepted route for majority of drugs but still faces challenges as compared to other routes. Transdermal route gives better control of blood level, reduce systemic toxicity, it protects drug from the hepatic first pass metabolism, better patient compliance and can be a potential option for oral dosage forms. Human skin is the important target site for the application of drug especially in the treatment of local disease. Penetration enhancement with special formulation approaches is mainly based on the usage of colloidal carriers [1]. Colloidal carrier have distinct advantages over conventional drug delivery as it act as drug containing reservoirs, modification of the particle composition or surface can adjust the release rate to the target site [2]. These carriers accumulate in stratum corneum or other upper skin layers and are not expected to penetrate into viable skin. The penetration enhancement is the most critical factor in the transdermal drug delivery [3]. Hence it is necessary to increase the flux through skin membrane by using different approaches of penetration enhancement. Vesicular systems have been widely studied as vehicles for dermal and transdermal drug delivery. A number of vesicles systems such as liposomes, niosomes, ethosomes, emulsomes and transfersomes have been developed. The vesicular carrier such as niosomes has distinct advantage over conventional dosage forms because these particles can act as drug reservoir [1-2].

Compounds having high molecular weight cannot cross skin, need some amendment owing to the availability of novel methods that might enhance the transport of large molecular weight compounds into or through the skin. Vesicular systems especially niosomes and liposomes are well established system for the transdermal route but the stability is the major problem of these carriers. Liposome exhibits some difficulties such as instability of aqueous dispersions on storage and the leakage of the encapsulated drugs. Also the high cost of synthetic phospholipids.

An alternative approach i.e. niosomes that overcomes several of these problems associated with liposomes [3]. Non-ionic surfactant vesicles obtained on hydration of synthetic non-ionic surfactants, with or

without incorporation of cholesterol or other lipid [4]. But the proniosomes are more advantageous than nonionic surfactant vesicles i.e., niosomes, in terms of physical stability such as aggregation, fusion and leaking and provide additional convenience in transportation, distribution, storage, and dosing [5]. Proniosomes encloses both hydrophilic and lipophilic drugs. Proniosomes reduce the toxicity related to drug because of their non-ionic nature of surfactant [6].

Neomycin Sulphate is a bactericidal aminoglycoside antibiotics (or antibacterial agent) is categorized as a BCS class-III i.e. high solubility and poor permeability, and generally it used as topical agent in skin infection. Physicochemical properties of NS like highly polar nature and high molecular weight (908.87D). Poor skin permeability (<3%) of Neomycin sulphate reduces its deeper penetration in skin [7]. That aminoglycoside antibiotic works by binding to the bacterial 30S ribosomal subunit, causing misreading of t-RNA, leaving the bacterium unable to synthesize proteins vital to its growth [8-10]. In present investigation proniosomal gel of NS was prepared by using span 60 and lecithin to increase the skin penetration.


### MATERIALS AND METHODS

#### Materials

Neomycin Sulphate was obtained as a gift sample from Encube Ethical Pvt. Ltd, Goa. Span 60, Cholesterol and carbopol 934 were purchased from Loba Chemicals, Mumbai. Lecithin was purchased from Research lab Mumbai. All other ingredients were of analytical grade.

#### Formulation of proniosomal gel

Proniosomes were prepared by the modified literature method reported by Fang *et al.* 2001 [11]. Proniosomes prepared by coacervation phase separation method. Precisely weighed amount of drug, surfactant, cholesterol, lecithin, and organic solvent (Ethanol) taken in wide mouth container. After mixing all ingredients, the open end of glass tube was covered with a lid to prevent loss of solvent from it and warmed on water bath at 60-70 °C for about 10 min, until the surfactants were dissolved completely. Then aqueous




Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



FORMULATION AND IN VITRO

journal.innovareacademic.in/index.php/ijap/article/view/30614

Innovare Home Our Journals Subscription Register Login



**International Journal of Applied Pharmaceutics**  
ISSN 0975-7058

Home About Current Archives Submissions Editorial Board Instructions To Authors Contact Us Search

HOME / ARCHIVES / VOL.11, ISSUE 2 (MAY-APRIL 2019) / Original Article(s)

## FORMULATION AND IN VITRO, IN VIVO EVALUATION OF PRONIOSOMAL GEL OF NEOMYCIN SULPHATE

**AMOL SHETE**  
Department of Pharmaceutics, Krishna Institute of Pharmacy of Krishna Institute of Medical Sciences Deemed To Be University, Karad-437012, India

**PRIYANKA THORAT**  
Research Group, Department of Pharmaceutics and Quality Assurance, Shree Santkrupa College of Pharmacy, Ghogaon, Karad-437014, India

**RAJENDRA DOIDAD**  
Department of Pharmaceutics, Krishna Institute of Pharmacy of Krishna Institute of Medical Sciences Deemed To Be


VIEW ABSTRACT  
PDF  
DOWNLOAD PDF  
HTML

PUBLISHED:

Type here to search

27°C Sunny 12:40 03-10-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



# Optimization of Thiazolidone Scaffolds Using Pocket Modeling for Development of Potential Secretory System Inhibitors of *Mycobacterium tuberculosis*

## *Mycobacterium tuberculosis*'in Potansiyel Sekreter Sistem İnhibitörleri Olarak Thiazolidone İskelelerinin Optimizasyonu

Shivratna V. KHARE<sup>1</sup>, Sujata P. CHOUDHARI<sup>2</sup>, Siddharth P. PHALLE<sup>1</sup>, Santosh S. KUMBHAR<sup>1</sup>, Prafulla B. CHOUDHARI<sup>1\*</sup>, Sambhaji R. MASAL<sup>1</sup>, Aakash K. PATIL<sup>1</sup>, Rakesh P. DHAVAL<sup>3</sup>, Durgacharan A. BHAGWAT<sup>3</sup>, Atul M. KADAM<sup>4</sup>

<sup>1</sup>Bharati Vidyapeeth College of Pharmacy, Department of Pharmaceutical Chemistry, Computational Chemistry Research Lab, Kolhapur, India

<sup>2</sup>Sarojini College of Pharmacy, Department of Pharmaceutical Analysis, Kolhapur, India

<sup>3</sup>Bharati Vidyapeeth College of Pharmacy, Department of Pharmaceutics, Kolhapur, India

<sup>4</sup>Shree Santkrupa College of Pharmacy, Department of Pharmaceutics, Ghogaon, India

### ABSTRACT

**Objectives:** *Mycobacterium tuberculosis* is the causative organism of tuberculosis, which is the most lethal disease after cancer in the current decade. The development of multidrug and broadly drug-resistant strains is making the problem of tuberculosis more and more critical. In the last 40 years, only one molecule has been added to the treatment regimen. Generally, drug design and development programs target proteins whose function is known to be essential to the bacterial cell. *M. tuberculosis* possesses specialized protein export systems like the SecA2 export pathway and ESX pathways.

**Materials and Methods:** In the present communication, rational development of an antimycobacterial agent's targeting protein export system was carried out by integrating pocket modeling and virtual analysis.

**Results:** The 23 identified potential lead compounds were synthesized, characterized by physicochemical and spectroscopic methods like infrared and nuclear magnetic resonance spectroscopy, and further screened for antimycobacterial activity using isoniazid as standard. All the designed compounds showed profound antimycobacterial activity.

**Conclusion:** We found that Q30, M9, M26, U8, and R26 molecules had significant desirable biological activity and specific interactions with Sec of mycobacteria. Further optimization of these leads is necessary for the development of potential antimycobacterial drug candidates with fewer side effects.

**Key words:** *Mycobacterium tuberculosis*, Sec, ESX, docking, antimycobacterial, multidrug resistant, pocket modeling

### ÖZ

**Amaç:** *Mycobacterium tuberculosis*, son on yılda kanserden sonra en ölümcül hastalık olan tüberkülozun etkenidir. Çoklu ilaç ve ilaca dirençli suşların gelişimi, tüberküloz problemini daha da kritik kılmaktadır. Son 40 yılda, tedavi rejimine sadece bir molekül eklenmiştir. Genel olarak ilaç tasarımı ve geliştirme programları, bakteri hücresi fonksiyonunda önemli olduğu bilinen proteinleri hedeflemektedir. *M. tuberculosis*, SecA2 ve ESX gibi özel protein ihracat sistemlerine sahiptir.


**Gereç ve Yöntemler:** Bu çalışmada, protein atım sistemini hedefleyen antimikobakteriyel bir bileşiğin rasyonel geliştirilmesi entegre cep modelleme ve sanal analiz kullanılarak gerçekleştirilmiştir.

\*Correspondence: E-mail: prafulla12@gmail.com - prafulla.choudhary@bharatividyapeeth.edu ORCID-ID: orcid.org/0000-0002-9137-3983

Received: 10/02/2018, Accepted: 22/03/2018

Turk J Pharm Sci, Published by Galenos Publishing House.



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Turkish Journal of Pharmaceut... x +

http://tjps.santkrupa.ac.in/archive-detail/article-preview/optimization-of-thiazolidone-scaffolds-using-pocket/26326

**TURKISH JOURNAL OF PHARMACEUTICAL SCIENCES**  
 Archive | Author Index | Turkish Pharmacists' Association, Academy of Pharmacy Web Page  
 E-ISSN: 2148-0247

About Us - Aims and Scope Editorial Board Advisory Board Instructions to Authors - Contact Us

**Original Article**

Optimization of Thiazolidone Scaffolds Using Pocket Modeling for Development of Potential Secretory System Inhibitors of *Mycobacterium tuberculosis*

10.4274/tjps.galenos.2019.12099

Shrivatna V. KHARE, Sujata P. CHOUDHARI, Siddhant R. PHALLE, Santosh S. KUMBHAR, Pratik B. CHOUDHARI, Sambhaj R. MASAL, Aakash K. PATIL, Rakesh P. DHAVALÉ, Durgacharan A. BHAGWAT, Anil M. KADAM

Turk J Pharm Sci 2019; 16(2): 196-205  
 PMID: 32454714

Received Date: 30.02.2018  
 Accepted Date: 23.03.2018

Abstract Full Text Video Image References 1319 14

Download PDF  
 How to Cite  
 Sitara  
 Send E-mail to Author  
 Share E-mail  
 Most Viewed Articles  
 Most Downloaded Articles  
 CC BY-NC-ND 4.0

Type here to search

27°C Sunny 12:42 03-03-2024



**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara

**Research Article**

**Analytical method development and validation of Valacyclovir to estimate from its formulation.**

**P. D. Lade, S. V. Patil, S. S. Kadam, O. B. Tipugade, P. G. Nakhare**

Research Group, Department of Pharmaceutics and Quality Assurance, Shree Santkrupa College of Pharmacy, Ghogaon, Karad-415111, Maharashtra, India.

Received 18 February 2019; received in revised form 08 March 2019; accepted 19 March 2019

\*Corresponding author E-mail address: [pdlade@gmail.com](mailto:pdlade@gmail.com)


**ABSTRACT**

A simple, rapid and sensitive method has been developed for the quantitative estimation of Valacyclovir hydrochloride in bulk and tablet. The zero order method was developed. The wavelength 254 nm was selected for the concentration range 5-25µg/ml. The accuracy of the method was assessed by recovery studies and was found to be 98.40±0.623. The method was statistically validated for the linearity, precision, accuracy repeatability, LOD, LOQ and ruggedness. The method was successfully applied for routine analysis of this drug in bulk and formulations. The method was validated as per ICH guidelines.

**KEYWORDS**

Valacyclovir, method development, validation.



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

3086

Analytical method developmen...  
 jcp:humanjournals.com/analytical-method-development-and-validation-of-valacyclovir-to-estimate-from-its-formulation/

**JCPR** Journal of Current Pharma Research  
 (An Official Publication of Human Journals)  
 An International Peer Reviewed Journal For Pharmacy, Medical & Biological Science  
 DOI: 10.25166 CODEN: JCPRD6 NLM ID: 101744065

ISSN Print: 2280-7834  
 Online: 2750-7847  
**HUMAN**  
 SJIF Impact Factor: 8.332

Home About Us Instructions to Author Publication Fee Manuscript Submission Archives Contact Us

**Analytical method development and validation of Valacyclovir to estimate from its formulation.**


11 FEBRUARY 2021  
 DOWNLOAD ARTICLE HERE: 26.P. D. Lode, S. V. Patil, S. S. Kadam, O. B. Tisugade, P. G. Nakhare (872.92 kB)  
 312 total views

Pharmaceutical Co-crystals: Modern solubility enhancement approach based on crystal engineering...  
 Synthesis, Characterization and Anticoagulant Activity of New Nitrogen substituted Chalcones

Search... Search  
 SJIF IMPACT FACTOR 8.332

27°C Sunny 12:52 03-02-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



Research Article

**Design, Development and Characterization of Fast Dissolving Oral Film of Clonazepam**

GOVIND G NIKAM, JAMEEL AHMED S MULLA\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Tal: Karad, Dist: Satara, Maharashtra - 415111

**ARTICLE DETAILS**

*Article history:*

Received on 18 February 2020

Modified on 12 March 2020

Accepted on 17 March 2020

*Keywords:*

Clonazepam,

Epilepsy,

Fast Dissolving Oral Film,

Solvent Casting,

HPMC E15.

**ABSTRACT**

The present study proposed to prepare fast dissolving oral film containing clonazepam for the treatment of epilepsy. HPMC E15 and PEG 400 were used as film forming agent and plasticizer, respectively. Solvent casting method was used to prepare Clonazepam loaded fast dissolving oral films. The prepared films were characterized for weight variation, thickness, percent elongation, tensile strength, folding endurance, moisture content, content uniformity, surface pH and swelling index. The DSC and FTIR Spectra revealed that drug was compatible with the polymer. The prepared oral films were opaque in nature having good folding endurance. Shows the rapid release of drug in oral cavity. Drug release by diffusion (93.44 %) and by dissolution (98.32%) after 5 minute. All nine batches are rapid release of drug after film contact with saliva.

© KESS All rights reserved

**INTRODUCTION**

According to US FDA, fast dissolving oral film is a sort of thin, flexible and non-friable polymeric film containing one or more dispersed active pharmaceutical ingredients. Fast dissolving oral film is intended to be kept in the buccal region for rapid disintegration or dissolution in the saliva before swallowing for drug delivery into the gastrointestinal tract [1].

In the last few decades, fast dissolving oral film has drawn much attention of researchers because of its distinct advantages over the other fast dissolving dosage forms. Fast dissolving oral film can be readily wetted and dissolved by saliva quickly without drinking water or chewing, allowing pediatric, geriatric or bedridden patients who have troubles in swallowing medicine [2, 3].

Fast dissolving oral film as a buccal drug delivery system, can enhance bioavailability by avoiding first-pass metabolism as well; as a result, patients can absorb drugs very fast through the rich vasculature of buccal region. For some special medicines which are vulnerable in gastrointestinal tract environment or have stomach irritation, it is also a good choice [4-7].

Epilepsy is the most common chronic brain disease and affects people of all ages. Nearly 50 million people around the world have epilepsy, which makes it one of the most common neurological diseases worldwide. Around 80% of people having epilepsy live in countries with lower and middle-income. It is characterized by recurrent seizures, which are short episodes of involuntary movement that may involve one part of the body (partial) or the whole body (generalized) and are sometime accompanied by loss of consciousness, control of bowel or bladder function [8].

Clonazepam (CNZ), a chlorinated derivative of nitrazepam, is an anticonvulsant benzodiazepine widely used in the treatment of epilepsy. It is also effective in the management of some types of neuralgia [9].

It is also used in myoclonus and associated abnormal movements, also for the treatment of panic disorders [10].

Moreover, it has recently been shown its efficacy also in the therapy of the burning mouth syndrome (BMS), pathology characterized by a painful burning sensation and/or other dysesthesias of the oral mucosa [11].


The present study intended to prepare fast dissolving oral film containing clonazepam for the treatment of epilepsy.

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

48



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Design, development and char: X +

ipdd.in/design-development-and-characterization-of-fast-dissolving-oral-film-of-clonazepam



Indian Journal of  
Novel Drug Delivery

ISSN 0975-5500



Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 12 [← Back to Issue](#)

January-March 2020

Design, development and characterization of fast dissolving oral film of clonazepam

Govind G Nikam, Jameel Ahmed S Mulla

[Download PDF](#)


**Abstract:**  
The present study proposed to prepare fast dissolving oral film containing clonazepam for the treatment of epilepsy. HPMC E15 and PEG 400 were used as film forming agent and plasticizer, respectively. Solvent casting method was used to prepare Clonazepam loaded fast dissolving oral films. The prepared films were characterized for weight variation, thickness, percent elongation, tensile strength, folding endurance, moisture content, content uniformity, surface pH and swelling index. The DSC and FTIR Spectra revealed that drug was compatible with the polymers. The prepared oral film shows superior in vitro release profile.

Type here to search

EUR/USD 0.87%

12:54  
23-10-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara





## A NOVEL CORTICOSTEROID CUBOSOMES – FOR OCULAR DRUG DELIVERY

Snehal Shashaikant Chakorkar<sup>1</sup>, Jameel Ahmed S. Mulla<sup>2</sup>

<sup>1</sup>Shree Vithal Education and Research Institutes College of Pharmacy, Pandharpur, Maharashtra, India.

<sup>2</sup>Shree Santkrupa College of Pharmacy, Ghogaon, Maharashtra, India.

### ARTICLE INFO

#### Article history

Received 14/05/2020

Available online

30/06/2020

#### Keywords

Cubosomes,  
Corticosteroids,  
Nanoparticle and Ocular Drug  
Delivery.

### ABSTRACT

Corticosteroid containing ocular formulation is very exigent tasks faced by the pharmaceutical industry while designing drugs acting on various ocular related diseases. Because ocular drug delivery faces various limitations like complex ocular anatomy and poor ocular bioavailability of drugs due to the high turnover rate of tears, lower corneal permeation, and rapid nasolacrimal drainage, and most important is irritation to the eye caused due to corticosteroid structure. The preferred dosage form for ocular delivery of corticosteroid is a solution or the ointment, but to sustain the level of drug at the target site during therapy is not possible so it is necessary to formulate novel drug delivery techniques. A large number of novel carrier drug delivery systems systems have been developed to overcome the above problems but among them, cubosomal drug delivery is a safe and effective technique for corticosteroid ocular drug delivery. Cubosomes are distinct, sub-micron; self assembles liquid crystalline particles having honeycomb (cavernous) structure which separate two internal aqueous channels and large interfacial area; having particle size ranges from 10-500 nm in diameter. This review briefly describes the ocular administration of corticosteroids with the cubosomal drug delivery system along with various drugs studied in ocular drug delivery, a method used for the preparation of Ocular cubosomes and its evaluation parameter. The benefits of cubosomal drug delivery will likely be applied widely in all treatment, diagnostic, and research aspects of ophthalmology in the future.

### Corresponding author

**Snehal Shashikant Chakorkar**

Shree Vithal Education and Research Institutes College of Pharmacy,

pandharpur, Maharashtra, India,

Pin code-413304

snehalchakorkar@gmail.com

Please cite this article in press as **Snehal Shashikant Chakorkar et al. A Novel Corticosteroid Cubosomes – for Ocular Drug Delivery. Indo American Journal of Pharmaceutical Research. 2020; 10(06).**

Copy right © 2020 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

[www.iajpr.com](http://www.iajpr.com)



**Principal**

**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

A NOVEL CORTICOSTEROID C...

https://www.zenodo.org/records/3503714

zenodo Search records Communities My dashboard Log in Sign up

INDO-AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH

Published June 30, 2020 | Version v1 Internal article Open

## A NOVEL CORTICOSTEROID CUBOSOMES – FOR OCULAR DRUG DELIVERY

Shetal Shashikant Chaturkar<sup>1</sup>, Jameel Ahmed S. Mulla<sup>2</sup><sup>1</sup>

Show affiliation

Corticosteroid containing ocular formulation is very exigent tasks faced by the pharmaceutical industry while designing drugs acting on various ocular related diseases. Because ocular drug delivery faces various limitations like complex ocular anatomy and poor ocular bioavailability of drugs due to the high turnover rate of tears, lower corneal permeation, and rapid nasolacrimal drainage, and most important is irritation to the eye caused due to corticosteroid structure. The preferred dosage form for ocular delivery of corticosteroid is a solution or the ointment, but to sustain the level of drug at the target site during therapy is not possible so it is necessary to formulate novel drug delivery techniques. A large number of novel carrier drug delivery systems systems have been developed to overcome the above problems but among them, cubosomal drug delivery is a safe and effective technique for corticosteroid ocular drug delivery. Cubosomes are distinct, sub-micron, self assemblies liquid crystalline particles having honeycomb (bicontinuous) structure which separate two internal aqueous channels and large interfacial area, having particle size ranges from 10-500 nm in diameter. This review briefly describes the ocular administration of corticosteroids with the cubosomal drug-delivery system along with various drugs studied in ocular drug delivery, a method used for the preparation of Ocular cubosomes and its evaluation parameter. The benefits of cubosomal drug delivery will likely be applied widely in all treatment, diagnostic, and research aspects of ophthalmology in the future.

141 VIEWS 309 DOWNLOADS Show more details


Versions

Version v1 June 30, 2020  
10.5281/zenodo.3503714

Cite all versions? You can cite all versions by using the DOI 10.5281/zenodo.3503714. This DOI represents all versions, and will always resolve to the latest one. Read more.

Type here to search 28°C Sunny 12:57 30-06-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## Research Article

**Formulation and Evaluation of Teneiglipitin-Loaded Mucoadhesive Microspheres**JAMEEL AHMED S MULLA<sup>1\*</sup>, VIJAYANAND R ARALELIMATH<sup>2</sup>, OMKAR TIPUGADE<sup>1</sup>, SHREYASI S SHINDE<sup>1</sup>, NISHA G TETGURE<sup>1</sup>, AYESHA A MULLA<sup>1</sup>, DINESH D GAVALI<sup>1</sup><sup>1</sup> Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, 415111, Maharashtra, India<sup>2</sup> Department of Pharmacology, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, 415111, Maharashtra, India**ARTICLE DETAILS***Article history:*

Received on 26 November 2020

Modified on 24 December 2020

Accepted on 28 December 2020

*Keywords:*Diabetes,  
Teneiglipitin,  
Mucoadhesive,  
Microspheres,  
In Vitro Release Study**ABSTRACT**

The mucoadhesive microspheres are one of the most promising novel techniques for drug delivery. Mucoadhesive systems provide a sustained drug release method, enhancing drug absorption in a site-specific manner. This study aims to prepare teneiglipitin mucoadhesive microspheres to increase the residence time in the gastric and offers control release. The teneiglipitin mucoadhesive microspheres are prepared by ionotropic external gelation technique using sodium alginate, HPMC K4, HPMC K15, HPMC K100, xanthan gum, guar gum, and carbopol 934. All mucoadhesive microspheres after preparation were characterized for percentage yield, particle size, drug entrapment efficiency, mucoadhesive test, and in vitro release. The results obtained are differed depending on the concentration of polymers and ratios of drug to polymers.

© KESS All rights reserved

**INTRODUCTION**

Diabetes is a chronic disease that occurs when the pancreas produces insulin inadequately or when the body cannot use the insulin produced by it efficiently. Insulin is a peptide hormone that controls the amount of glucose in the blood, commonly known as blood sugar. Hyperglycaemia, or raised blood sugar, is a common effect of uncontrolled diabetes. Unchecked diabetes, over time, leads to potential complications in various organ systems of the body, which includes heart ailments, kidney damage, and nerve damage. According to The World Health Organization (WHO), about 8.5% of adults aged 18 years and above had diabetes in 2014. During 2016, diabetes was the direct cause of 1.6 million deaths, and in the year 2012, high blood glucose was the leading cause of 2.2 million deaths additionally [1-2]. Nearly half of all deaths attributable to high blood glucose occur before 70 years of age. WHO estimated the diabetes was the seventh leading cause of death in 2016 [2].

Dipeptidyl peptidase-4 (DPP-4) inhibitors are a comparatively new form of oral diabetes drugs. Also identified as gliptins, they are generally prescribed for people with type-2 diabetes who do not respond well to medications such as metformin and sulphonylureas. DPP-4 is an enzyme that destroys the hormone incretin. DPP-4 inhibitors work by blocking the activity of DPP-4. Incretins help the body produce more insulin only when needed and reduce the liver's amount of glucose when it is not required. These hormones are released during the whole day, and levels are increased at mealtimes [3].

Teneiglipitin is one of the newly approved gliptins and is effective in treating type-2 diabetes. It is hypothesized that an oral formulation that could substantially retain in the gastrointestinal tract (GIT) and release the drug in a controlled manner could be highly effective than a single dose conventional dosage form. In this context, mucoadhesive drug delivery systems adhere to certain gastrointestinal segments and would offer various advantages.

The present study aims to prepare mucoadhesive microspheres of teneiglipitin to prolong the residence time in the gastric and provide control release.

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

222



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Formulation and evaluation of...

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 12

October-December 2020

Formulation and evaluation of teneligliptin-loaded mucoadhesive microspheres

Janghel Abinad S Mulla, Vijayamand R Aralelimath, Omkar Tipogade, Shreyasi S Shinde, Nisha G Tetgire, Ayesha A Mulla, Dinesh D Gavali


Download PDF

**Abstract:**  
The mucoadhesive microspheres are one of the most promising novel techniques for drug delivery. Mucoadhesive systems provide a sustained drug release method, enhancing drug absorption in a site-specific manner. This study aims to prepare teneligliptin mucoadhesive microspheres to increase the residence time in the gastric and offers control release. The teneligliptin mucoadhesive microspheres are prepared by ionicotropic external gelation technique using sodium alginate, HPMC K4, HPMC K15, HPMC K100, xanthan gum, guar

Type here to search

28°C Sunny 12:59 23-10-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



## Characterization, antioxidant, antimicrobial and cytotoxic activities of green synthesized silver and iron nanoparticles using alcoholic *Blumea eriantha* DC plant extract

Rohankumar R. Chavan<sup>a</sup>, Somnath D. Bhinge<sup>a,\*</sup>, Mangesh A. Bhutkar<sup>b</sup>, Dheeraj S. Randive<sup>b</sup>, Ganesh H. Wadkar<sup>c</sup>, Sachin S. Todkar<sup>d</sup>, Mukund N. Urade<sup>d</sup>

<sup>a</sup> Department of Pharmaceutical Chemistry, Rajarambapu College of Pharmacy, Kasegaon, Maharashtra, 415 404, India

<sup>b</sup> Department of Pharmacokinetics, Rajarambapu College of Pharmacy, Kasegaon, Maharashtra, 415 404, India

<sup>c</sup> Department of Pharmacognosy, Rajarambapu College of Pharmacy, Kasegaon, Maharashtra, 415 404, India

<sup>d</sup> Department of Pharmacology, Shri Santkrupa College of Pharmacy, Ghogaon, Maharashtra, 415 111, India

<sup>e</sup> Department of Pharmaceutical Chemistry, Adarsh College of Pharmacy, Vili, Maharashtra, 415 311, India

### ARTICLE INFO

#### Keywords:

Antibacterial  
Antioxidant  
Antimicrobial  
Cytotoxicity  
MCF-7  
Apoptosis  
DAPI staining

### ABSTRACT

Silver and iron nanoparticles was synthesized, characterized and investigated for biological screening using alcoholic *Blumea eriantha* DC plant extract. Equal amount of plant extract was mixed and incubated with silver nitrate and ferric chloride to obtain silver and iron nanoparticles respectively. Formation of silver and iron nanoparticles was confirmed by using UV, FT-IR spectroscopy, SEM, X-ray diffraction, and TEM. The anti-oxidant, antimicrobial, cytotoxic activities of the synthesized nanoparticles were determined by using standard protocols. MCF-7 cells were treated with selected concentrations of extract and prepared nanoparticles. It was observed that the synthesized nanoparticles were spherical in shape with an average particle size of 50 nm. The results of the studies revealed that the synthesized nanoparticles exhibited effective antioxidant, antibacterial and cytotoxic activity. The cell viability was measured for 50 hours after the addition of selected concentration of nanoparticles. The inhibition rate of silver nanoparticles was observed to be 15.45, 20.25 and 28.16 % against concentration 25, 50 and 100  $\mu\text{g mL}^{-1}$  respectively. DPPH staining and Annexin V FITC assay results indicated that silver nanoparticle induce apoptosis in MCF-7 cells as compared with control. Therefore, it may open up a new avenue for anticancer therapies that needs further research.

### 1. Introduction

Dignified metal nanoparticles are becoming increasingly trendy, more than ever, due to their exceptional biocompatibility [1]. As a new material, silver nanoparticles (AgNPs) are considered as one of the most extensively used metallic nanoparticles in the era due to the wide physicochemical properties. Also their wide physicochemical properties raised much concern in the biomedical field, the pharmaceutical industry etc [1–3]. Moreover, AgNPs are in an alternative quite good agent to carry such effective drug moieties at the targeted area [4]. Iron is mainly plentiful and widely used elements on earth [5]. Silver and iron have been reported to exhibit a wide array of pharmacological activities since ancient times [6–10]. NPs are well known for their beneficial biomedical inhibitory properties such as anticancer,

antibacterial, antifungal, antiviral, etc. [11].

Being environmentally friendly, cost-effective and easily scalable, green synthesis is an excellent substitute for the electrochemical, chemical, radiation, photochemical, Langmuir–Blodgett and biological synthesis [12]. However, green synthesis have attracted merit importance due to its one-step, faster, cost-effective, nontoxic and eco-friendly synthetic approach [13]. In addition, earlier methods have already proved the successful synthesis of NPs using bacteria [14,15], plant extract [16,17], fungus [18], panchakavya [19] and cow milk [20]. The use of phytomedicine has become very popular with time owing to its medicinal properties [12]. In nanotechnology the plant extracts are not only a good way to make nanoparticles that are harmless and stable, but also a good way to reduce the use of toxic chemicals or powerful reducing agents.

\* Corresponding author at: Department of Pharmaceutical Chemistry, Rajarambapu College of Pharmacy, Kasegaon, Dist - Sangli, Maharashtra, 415404 India.

E-mail addresses: [rohankumar3102@gmail.com](mailto:rohankumar3102@gmail.com) (R.R. Chavan), [somu1245@yahoo.co.in](mailto:somu1245@yahoo.co.in) (S.D. Bhinge), [mangesh\\_bhutkar@rediffmail.com](mailto:mangesh_bhutkar@rediffmail.com) (M.A. Bhutkar), [randivedheeraj@gmail.com](mailto:randivedheeraj@gmail.com) (D.S. Randive), [ganeshpharma77@gmail.com](mailto:ganeshpharma77@gmail.com) (G.H. Wadkar), [sachintodkar007@gmail.com](mailto:sachintodkar007@gmail.com) (S.S. Todkar), [mukundurade@gmail.com](mailto:mukundurade@gmail.com) (M.N. Urade).


<https://doi.org/10.1016/j.mtcomm.2020.101320>

Received 19 March 2020; Received in revised form 17 May 2020; Accepted 4 June 2020

Available online 13 June 2020

2382-4928/© 2020 Elsevier Ltd. All rights reserved.



  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Characteristics antioxidant

sciencedirect.com/science/article/doi/10.1016/j.matcom.2020.101323

ScienceDirect Journals & Books Search... Register Sign In

Access through your institution Purchase PDF

materialstoday COMMUNICATIONS

Volume 24, September 2020, 101323

Characterization, antioxidant, antimicrobial and cytotoxic activities of green synthesized silver and iron nanoparticles using alcoholic *Blumea eriantha* DC plant extract

Rubankumar R. Chavan<sup>a</sup>, Sampath D. Bhatia<sup>a</sup>, Manoj A. Shukla<sup>a</sup>, Dheeraj S. Bendre<sup>a</sup>, Ganesh H. Wadkar<sup>a</sup>, Sachin S. Todkar<sup>a</sup>, Mukund N. Urade<sup>a</sup>

Recommended articles

Antifungal and antibacterial activity of densely dispersed silver nanospheres...  
Journal de Mycologie Médicale, Volume 24, Issue ...  
M. Khatami, ... R. Singh

Super-paramagnetic iron oxide nanoparticles (SPIONs): Greener...  
Journal of Cleaner Production, Volume 248, 2020 ...  
Mehrdad Khatami, ... Arif Ullah Khan

Mikania mikromidia leaf extract mediated biogenic synthesis of magnetic iron...  
Materials Today: Proceedings, Volume 42, Part 2, 3 ...  
Anayuki Biswal, ... Sarvesh Lathwasia Nakhum

Show 3 more articles

FEEDBACK

Type here to search 28°C Sunny 13:01 03-10-2024



**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



Review Article

**Microemulsion Based Hydrogel Formulation for Topical Drug Delivery - A Concise Review**

JAMEEL AHMED S MULLA\* AND BIRADEV S KARANDE

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Maharashtra, India - 415111

**ARTICLE DETAILS**

*Article history:*

Received on 15 April 2021

Modified on 22 May 2021

Accepted on 25 May 2021

*Keywords:*

Microemulsion,  
Microemulsion-based  
Hydrogel,  
Ternary Phase Diagram,  
Preparation,  
Characterization.

**ABSTRACT**

A hydrogel is a network of water-insoluble polymer chains that can also be found as a colloidal gel with water as the dispersion medium. Hydrogels are natural or manufactured polymers that are superabsorbent (they can hold over 99 percent water). Topical medicines are utilized for localized effects at the application site due to medication penetration into the deeper layers of the skin or mucous membranes. Microemulsions are thermodynamically stable, fluid, transparent (or translucent) colloidal dispersions made up of an oil phase, aqueous phase, surfactant, and co-surfactant in appropriate ratios that form a single optically isotropic solution with droplet diameters typically ranging from 10 to 100 nanometers. Transparency, low viscosity, and, most importantly, thermodynamic stability and capacity to form spontaneously separate micro-emulsions from conventional emulsions. As a topical medication delivery system, micro-emulsions provide a number of advantages over standard creams, gels, and solutions. The Hydrogel technology based on microemulsion will be able to sustain therapeutic concentration at the site of action while also increasing bioavailability. This review focuses on the method of preparation, characterization, evaluation, and stability investigations of microemulsion-based hydrogel.

© KESS All rights reserved

**INTRODUCTION**

Pharmaceutical research nowadays is focused on finding appropriate drug delivery systems to meet the therapeutic demands of patients. The discovery of new chemical entities should coincide with the development of novel drug delivery systems, as the majority of compounds identified were hydrophobic in nature. Microemulsion has gained significance in the effective delivery of hydrophobic medications as part of the push to create innovative delivery strategies [1]. In recent years, the creation of so-called "intelligent" or "smart" hydrogels has gotten a lot of attention in soft matter research. Hydrogels having a high water content that can undergo reversible phase transitions in response to changes in temperature, pH, magnetic field, light, solvent, and ionic strength are known as "intelligent" hydrogels.

Hydrogels made of N-isopropyl acrylamide (NiPAAm) are known to change volume and transparency when heated above 32°C, the gel's lower critical solution temperature (LCST) [2]. Microemulsion is a transparent emulsion that is frequently referred to as O/W or W/O emulsion.

Product with droplet sizes ranging from 10 to 100 nm and little tendency to agglomerate. It is made up of a proper ratio of oil phase, surfactant, co-surfactant, and aqueous phase. Transparency, optical isotropy, low viscosity, and thermodynamic stability are among microemulsion's physicochemical features. Topical drug administration appears to be a viable route of drug administration. Several mechanisms have been postulated to explain the benefits of microemulsions for drug administration in the topical and dermal areas. First, due to the huge amount of a drug integrated in the formulation, the thermodynamics towards the skin are boosted. Second, the drug's higher thermodynamic activity could help it partition into the skin. Third, by serving as permeation enhancers, the microemulsion components may diminish the

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com



**Principal**

**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Microemulsion based hydrogel

ipdd.in/microemulsion-based-hydrogel-formulation-for-topical-drug-delivery-A-concise-review/



Indian Journal of  
Novel Drug Delivery

ISSN 0975-5500



Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 13 [← Back to Issue](#)

April-June 2021


Microemulsion based hydrogel formulation for topical drug delivery - A concise review

Jameel Ahmed S Mulla, Biradev S Karande

[Download PDF](#)

**Abstract:**  
A hydrogel is a network of water-insoluble polymer chains that can also be found as a colloidal gel with water as the dispersion medium. Hydrogels are natural or manufactured polymers that are superabsorbent (they can hold over 99 percent water). Topical medicines are utilized for localized effects at the application site due to medication penetration into the deeper layers of the skin or mucous membranes. Microemulsions are thermodynamically stable, fluid, transparent (or translucent) colloidal dispersions made up of an oil phase.

Type here to search 28°C Sunny 13:03 23-02-2024



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Contents lists available at ScienceDirect

Journal of Drug Delivery Science and Technology

journal homepage: [www.elsevier.com/locate/jddst](http://www.elsevier.com/locate/jddst)

Research paper

## Effect of polymers and process parameters in augmenting the compactability and dissolution behaviour of oxcarbazepine spherical agglomerates

Sandip Honmane<sup>a,d,\*</sup>, Atul Kadam<sup>b,d</sup>, Sujata Choudhari<sup>c</sup>, Raviraj Patil<sup>d</sup>,  
Siddique Akber Ansari<sup>e</sup>, Vinod Gaikwad<sup>f</sup>

<sup>a</sup> Department of Pharmaceutics, Anasaheb Dange College of B. Pharmacy, Ashta, 416301, Maharashtra, India

<sup>b</sup> Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, 415111, Maharashtra, India

<sup>c</sup> Department of Pharmaceutical Analysis, Sarojini College of Pharmacy Kolhapur, 416013, Maharashtra, India

<sup>d</sup> Department of Pharmaceutics, Ashokrao More College of Pharmacy, Feth-Vadgaon, 416112, Maharashtra, India

<sup>e</sup> Department of Pharmaceutical Chemistry, College of Pharmacy, King Saud University, Riyadh, P.O. Box 2454, Riyadh 11452, Saudi Arabia

<sup>f</sup> Department of Pharmaceutics, BVDCU Poona College of Pharmacy, Pune, 411038, Maharashtra, India



## ARTICLE INFO

## Keywords:

Spherical crystallization

Agglomerate

Micrometric

Solubility

Compactability

## ABSTRACT

The hypothesis for the present work is proposed to explore the spherical crystallization technique for improving the micromeritics, compactability, and solubility characteristics of oxcarbazepine (OXZ), an anticonvulsant drug. Agglomerates containing different polymers (PEG 6000 and PVP K30) and process parameters were investigated for the enhancement of overall physicochemical performance and dissolution. Water, dichloromethane, and chloroform were used as a poor solvent, good solvent, and bridging liquid, respectively. Pure OXZ and spherical agglomerates were characterized for several properties including Fourier transformation-infrared spectroscopy, Differential scanning calorimetry, Scanning electronic microscopy, X-ray powder diffraction analysis, micromeritics, solubility studies, and *in-vitro* drug release kinetics. From the results, a considerable improvement in drug solubility and micromeritics of agglomerates than pure OXZ was observed. Compressibility parameters assessed from the Heckel plot showed agglomerates with a polymer having a higher value of slope (k) and less MyP is accountable for plastic deformation in agglomerates. Prepared spherical agglomerates showed an enhancement in solubility and rate of dissolution, which might improve their bioavailability.

## 1. Introduction

The recent trend in the pharmaceutical industry is focused to deliver quality medicines developed through a pharmaco-economic approach. Tablet is well accepted solid dosage form by the patients for the oral administration of the drugs. Drugs with poor flowability and mechanical properties of crystal-like compressibility are unsuitable for direct compression process; and require wet granulation which is uneconomical, laborious, and tedious. Techniques used to improve flow properties of compression mixture comprised of extrusion-spheronization, melt solidification, melt granulation, melt extrusion, and spherical crystallization [1]. Spherical agglomeration (Crystallo-co-agglomeration) is a recent alternative technique wherein *in-situ* formation and agglomeration of crystals is carried out to prepare directly compressible

agglomerates [2–4]. This technique allows simultaneous crystallization and agglomeration in a single step, to transform the fine spherical crystals produced, directly into the more compactable and directly compressible form [5,6]. Apart from modifications in the primary and secondary properties of the particles, the Crystallo-co-agglomeration technique not only reduces the number of unit operations and processing costs but also improves the dissolution rate and bioavailability of the actives [7,8]. Moreover, during processing, modification of crystal habit with improved physicochemical (solubility, dissolution rate, and stability) and micromeritic properties (flowability, and compactability) [9] has emerged as a potential area of research in pharmaceutical manufacturing. Consequently, the directly compressible spherical agglomerates formed by this technique exhibited a considerable impact on the formulation of dosage forms [6,10].

\* Corresponding author. Department of Pharmaceutics, Anasaheb Dange College of B. Pharmacy, Ashta, 416 301, Maharashtra, India.  
E-mail address: [sandiphonmane@gmail.com](mailto:sandiphonmane@gmail.com) (S. Honmane).


<https://doi.org/10.1016/j.jddst.2021.102576>

Received 26 December 2020; Received in revised form 25 April 2021; Accepted 6 May 2021

Available online 11 June 2021

1773-2247/© 2021 Elsevier B.V. All rights reserved.



  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Effect of polymers and process... | ScienceDirect | Journals & Books | Search... | Register | Sign In

Access through your institution | Purchase PDF | Patient Access

Journal of Drug Delivery Science and Technology  
Volume 64, August 2021, 102578

Research paper

### Effect of polymers and process parameters in augmenting the compactability and dissolution behaviour of oxcarbazepine spherical agglomerates

Sandya Homare<sup>a,\*</sup>, Atul Kadam<sup>b,c</sup>, Sujata Chaudhari<sup>b</sup>, Ravini Patil<sup>d</sup>, Siddique Akbar Ansari<sup>e</sup>, Vinod Gaikwad<sup>f</sup>

Show more

Add to Mendeley | Show | Cite

Recommended articles


- Effect of USP induction ports and modified glass sampling apparatus on...  
Journal of Drug Delivery Science and Technology...  
Riyadh R. Mehrez, ... Almaraz R. Ponce
- Influence of modified induction port, modified DUSA assembly and device at...  
Journal of Drug Delivery Science and Technology...  
Riyadh R. Mehrez, ... Almaraz R. Ponce
- Effect of the cargo lipophilicity on powder micromeritics properties of...  
Journal of Drug Delivery Science and Technology...  
Cheng Ma, ... Chuanbin Wu

Show 3 more articles

FEEDBACK

28°C Sunny | 13:25 | 23-02-2024



  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



## Research Article

**Formulation and Characterization of Rapidly Dissolving Buccal Films of Montelukast Sodium**

ANKITA B HOGALE, JAMEEL AHMED S MULLA\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Maharashtra, India - 415111

**ARTICLE DETAILS***Article history:*

Received on 20 July 2021

Modified on 21 August 2021

Accepted on 25 August 2021

*Keywords:*Rapidly Dissolving Film,  
Montelukast Sodium,  
Solvent Casting Method,  
Buccal Film.**ABSTRACT**

In the event of chronic illnesses, fast-dissolving drug delivery systems have been created as an alternative to traditional dosage forms as an oral mode of drug delivery. Fast dissolving films are now favoured over traditional tablets and capsules for disguising the taste of bitter medications and increasing patient compliance. In this study, montelukast sodium-loaded fast dissolving oral films were made using HPMC K-100 and the solvent casting method. Disintegration time, thickness, tensile strength, percent elongation, folding endurance, moisture content, surface pH, content uniformity, swelling index, FTIR Spectroscopy, Differential Scanning Calorimetry, and an *in-vitro* dissolution investigation were all used to describe the prepared films. The chemical structure of montelukast sodium was preserved in the formulation, according to FTIR analysis. At the end of 5 minutes, the drug release was found to be between 90.75 and 99.14 percent.

© KESS All rights reserved

**INTRODUCTION**

In the late 1970s, rapid-dissolving drug-delivery systems were developed as an alternative to tablets, capsules, and syrups for juvenile, geriatric, bedridden, nauseated, or noncompliant patients who had difficulty ingesting typical oral solid-dosage forms. The technique, which was based on the technology of transdermal patches, is also known as oral thin films, fast dissolving films, mouth dissolving films, oro-dispersible films, quick disintegrating films, and melt in mouth dosage form [1,2]. Drug administration via rapidly dissolving buccal films has developed as a cutting-edge alternative to the usual pills, capsules, and liquids commonly used in prescription and over-the-counter drugs. Thin film strips are primarily designed for oral administration, with the user placing the strip on or under the tongue or along the inside of the cheek, and are similar in size, shape, and thickness to a postage stamp. When a film is placed on the tongue or buccal cavity, it dissolves quickly, sending the medication to the systemic circulation via breakdown [3,4]. Montelukast sodium is an anti-asthmatic medicine that works as a leukotriene receptor antagonist, highlighting the importance, optimal qualities, and many

aspects of mouth dissolving film formulation as a superior dosage form for treating asthma and improving patient compliance [5].

The primary goal of this study is to develop a fast dissolving film of montelukast sodium, an anti-asthmatic medicine, which will aid in the rapid beginning of action and increase the performance of the active pharmaceutical ingredient.

**MATERIALS AND METHODS****Materials**

Aarti Pharma in Mumbai provided the Montelukast Sodium. Loba Chemie, Mumbai, provided HPMC K100, PEG 600, and sodium lauryl sulphate. All of the other chemicals, excipients, and solvents utilized were obtained from reputable sources and are of laboratory and analytical quality.


**Formulation of Fast Dissolving Films of Montelukast Sodium**

Solvent casting was used to make the quickly dissolving films, with HPMC K-100 as the film former. With constant stirring on a magnetic stirrer, the calculated amount of polymer, HPMC K-100, was dissolved in one fourth amount of distilled water. The montelukast sodium was adequately dissolved in a suitable amount of ethanol and then added to the polymeric solution

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Formulation and characterizati... x +

jdidd.in/formulation-and-characterization-of-rapidly-dissolving-buccal-films-of-montelukast-sodium/



Indian Journal of  
Novel Drug Delivery

ISSN 0975-5500



Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 13 [← Back to Issue](#)

July-September 2021

**Formulation and characterization of rapidly dissolving buccal films of montelukast sodium**

Ankita B Hogale, Jameel Ahmed S Mullis


[Download PDF](#)

**Abstract:**  
In the event of chronic illnesses, fast-dissolving drug delivery systems have been created as an alternative to traditional dosage forms as an oral mode of drug delivery. Fast dissolving films are now favoured over traditional tablets and capsules for disguising the taste of bitter medications and increasing patient compliance. In this study, montelukast sodium-loaded fast dissolving oral films were made using HPMC K-100 and the solvent casting method. Disintegration time, thickness, tensile strength, percent elongation, folding

Type here to search

Freezing news 13:54 23-02-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

## Ecopharmacovigilance: Perspectives, concepts, applications, and relationships in modern context

Prachi Khamkar<sup>1\*</sup>, Debarshi Kar Mahapatra<sup>2</sup>, Atul Kadam<sup>3</sup>

<sup>1</sup>Department of Pharmaceutics, University of Mumbai, Mumbai 400020, Maharashtra, India

<sup>2</sup>Department of Pharmaceutical Chemistry, Dadasaheb Balpande College of Pharmacy, Nagpur 440037, Maharashtra, India

<sup>3</sup>Department of Pharmaceutics, Shree Santakrupa College of Pharmacy, Ghogaon, Shivaji University, Kolhapur 416012, Maharashtra, India

\*Corresponding author/authors complete details (E-mail and Telephone): Email ID: prachipkhamkar15@gmail.com, Ph. No: +916888199055

DOI: 10.5281/zenodo.4568961

Web of Science Researcher ID: AAG-6443-2021

ORCID ID: 0000-0002-8579-8858

### Abstract

Date Received: 04/01/2021  
Date Revised: 05/02/2021  
Date Accepted: 18/02/2021

#### Keywords

Ecopharmacovigilance,  
Pharmacovigilance,  
Pharmaceuticals, Waste,  
Environment, Removal

Environmental scientists have made great strides to regulate pharmaceutical waste. However, the monitoring of emerged environmental problems induced by drugs should attract further interest of pharmacy and pharmacovigilance scientists. Ecopharmacovigilance (EPV) as a kind of pharmacovigilance for the atmosphere is widely recognized as essential to minimize the environmental impact of pharmaceutical toxins. In efforts to answer the environmental issues created by medications, the constructive involvement of the pharmaceutical sector is essential. In a prioritized basis, EPV can target individual pharmaceuticals. For EPV deployment targeting pharmaceutical contamination, certain advice and management practice solutions are recommended. On administration certain drugs are metabolized throughout the bloodstream, being inert or becoming converted to metabolites, whereas others are excreted in the urine or liver and excreted in the sewage. The substances that are released into drainage can be processed into a number of chemicals by a sewage treatment process. The involvement of different drugs and their components has been found in the marine world, with the aquatic environment being the most researched to-date. Nanostructure materials have been around for a long time, and its interactions with biological processes have been discussed in various applications to enhance the understanding and importance of environmental and health effects. In order to regulate pharmaceutical residues in effluents, Technologies for sewage water management should be applied. In particular, the presence of pharmacy including pharmacovigilance professionals is also important for enhancing multidisciplinary collaboration.


An official publication of Global Pharmacovigilance Society; Published under licence of Creative Commons Attribution 4.0 International. COPYRIGHT © 2020 JPADR



### Introduction

The research of "Ecopharmacovigilance" (EPV) has become a recent subject of concern in the sense of the rise of pharmaceutical contamination. For the last 30 years, multinational institutions and the pharmaceutical industry have come to recognize that pharmaceutical drugs have a negative global effect on the climate. Although it is a global issue, like other environmental issues, pharmaceutical pollution more directly and seriously affects those living near production plants whose water and food sources are contaminated with waste pharmaceutical products (Figure 1) (Wang and Hu, 2014). With the continued rapid development of the global pharmaceutical industry, increasing attention has been given to environmental issues caused by pharmaceutical

pollutants. This is an increase in the consumption, occurrence, and persistence of pharmaceutical products in the environment with their diverse biological effects (Coettier *et al.*, 2007). While not entirely, although to a great degree, medicines were a blessing for mankind in curbing the diseases (Medhi and Sewal, 2012). In the developed world, pharmacovigilance was well accepted and practiced, but decades were absorbed by emerging areas of the globe, India is also currently starting a national initiative to track the adverse effects of medications (Gupta, 2010). These environmental pharmaceutical pollutants include excretion of pharmaceuticals after human and veterinary therapeutic use. This dominates the global input of pharmaceuticals into the environment and are a much more difficult source to control: adding to the direct release into the wastewater

**Principal**  
Dr. Ramling G. Patrakar  
Shree Santakrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Ecopharmacovigilance Persp...

ipadr.com/index.php/jpadr/article/view/13

Register Login

## Journal of Pharmacovigilance and Drug Research

Home About Contents About Previous Issues Guidelines Publication Ethics Events Sponsors Search

Announcements

Home > Archives > Vol 2 No 1 (2021): Journal of Pharmacovigilance and Drug Research > Reviews

### Ecopharmacovigilance: Perspectives, concepts, applications, and relationships in modern context

**Prachi Khamkar**  
University of Mumbai, Mumbai 400032, Maharashtra, India  
[prachi.khamkar@uom.ac.in](mailto:prachi.khamkar@uom.ac.in)

**Debarshi Kar Mahapatra**  
Dadasaheb Barsekar College of Pharmacy, Naggur 440037, Maharashtra, India

**Atul Kadam**  
Shree Santkrupa College of Pharmacy, Ghogaon, Shree University Kolhapur 416012, Maharashtra


DOI: <https://doi.org/10.53411/jpadr.2021.2.1.7>

Make a Submission

Information  
For Readers  
For Authors  
For Librarians

Keywords



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara





## Regulatory challenges and myths in pharmaceutical 3D printing

Although three-dimensional (3D) printing has been around for some time, many people still underestimate its capabilities. This article addresses the regulatory challenges faced by the pharmaceutical sector and aims to correct some of the most common myths surrounding the technology.

**I**MAGINE A FUTURE where patients with multiple disease conditions no longer have to take several tablets, numerous times per day. Instead, they can take one tablet containing all their required medications, once daily, thanks to 3D printing.

The process of 3D printing begins with a computer-aided format of a digital prototype of the product. The format is then sliced into horizontal layers that will form the shape of the digital file before being transferred into a 3D printer. Various excipients and drug combinations are then used to print the product with the aid of layering, transforming two-dimensional (2D) layers into a 3D product. Although the pharmaceutical industry


is a laggard in this area, it is now waking up to the possibilities of 3D printing and we have witnessed various developments whereby specific tailor-made medicines can be manufactured according to a patient's need.

### Where are we now?

The four most frequently used 3D printing machines in the pharmaceutical industry are:

1. Stereolithography (SLA)
2. Fused deposition modelling (FDM)
3. Selective laser sintering (SLS)
4. Binder jetting (BJ).



  
**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



Regulatory challenges and myths in pharmaceutical 3D printing

STAY UPDATED  
SUBSCRIBE TODAY!

actylis Custom GMP Manufactured Raw Materials

INNOVATE WITH GEA PHARMA AT ACHEMA.  
10-14 June 2024  
Frankfurt am Main, Germany.  
ACHEMA2024

Endosafe Trillium  
A Recombinant Cascade Respart (RCR)

By Atul Kadam (Shree Santkrupa College of Pharmacy, Prachi Khambur) (Shree Santkrupa College of Pharmacy)

18 February 2021

No comments yet

Although three-dimensional (3D) printing has been around for some time, many people still underestimate its capabilities. This article addresses the regulatory challenges faced by the pharmaceutical sector and aims to correct some of the most common myths surrounding the technology.



*[Handwritten Signature]*  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

**FORMULATION AND PROCESS VALIDATION OF  
CLARITHROMYCIN LOADED IMMEDIATE RELEASE TABLETS****Ketaki S. Shinde\*<sup>1</sup>, Dr. R. C. Doijad<sup>1</sup>, Dr. J. S. Mulla<sup>1</sup> and Sachin S. Mali<sup>2</sup>**<sup>1</sup>Department of Quality Assurance, Shree Santkrupa College of Pharmacy, Shivaji University, Ghogaon, India.<sup>2</sup>Department of Pharmaceutics, Y. D. Mane Institute of Pharmacy, Kagal, India.Article Received on  
17 January 2022,Revised on 06 Feb. 2022,  
Accepted on 27 Feb. 2022

DOI: 10.20959/wjpps20223-21006

**\*Corresponding Author****Ketaki S. Shinde**Department of Quality  
Assurance, Shree Santkrupa  
College of Pharmacy,  
Shivaji University,  
Ghogaon, India.**ABSTRACT**


The main aim of the present research work was to study concurrent process validation of immediate release tablet. If each step of production process is validated we can assure that the final product is of the best quality. Validation is best viewed as an impartment and integral part of cGMP. Validation is therefore one element of quality assurance programs associated with a particular process. Quality cannot be assured only by doing finished product testing and in-process monitoring but it should be built into the manufacturing process. So building of quality require a special attention to a few factors like selection of material, process design, control variables, in process control and finished product testing. In this study three initial

batch of clarithromycin tablet with same size, method, equipment and validation criteria were taken. The critical parameters involved in dry mixing, preparation of granulating agent, wet mixing, wet milling, drying, lubrication, and compression stages were identified and evaluated. Results obtained with this process validation data provides high degree of assurance that manufacturing process produces product meeting its predetermined specifications and quality attributes. It also provides documented evidence for the operation sequence of manufacturing process and to determine the critical parameters and variables in the process of manufacturing of the tablets. The output of process validation can be used to increase productivity, its consistent quality and decreasing the need for processing or market complaints.

**Principal**  
**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

WJPS | ABSTRACT

WJPS.COM/WJPS\_CONSOLE/ABSTRACT\_16/16135



**WORLD JOURNAL OF PHARMACY AND  
PHARMACEUTICAL SCIENCES**  
(An ISO 9001:2015 Certified International Journal)


An International Peer Reviewed Journal for Pharmaceutical and Medical Research and Technology

ISSN 2278-4367  
Impact Factor : 7.632  
ICV : 84.65

SAT, FEB 03 2024 | 3:49:50 PM

World Journal of Pharmacy and Pharmaceutics

[Login](#) / [Register](#)



---

HOME | ABOUT US | INSTRUCTION TO AUTHOR | CURRENT ISSUE | MANUSCRIPT SUBMISSION | TRACK YOUR ARTICLE | ARCHIVE | PROCESSING FEES | CONTACT US

**WJPS Citation** Total

	All	Since 2019
Citation	1450	3999
h-index	23	20
i10-index	134	54

### Abstract

**FORMULATION AND PROCESS VALIDATION OF CLARITHROMYCIN LOADED IMMEDIATE RELEASE TABLETS**

Katuki S. Shinde\*, Dr. R. C. Dojrad, Dr. J. S. Mulla and Sachin S. Mat

**ABSTRACT**

The main aim of the present research work was to study, formulate process validation of immediate release tablet. If each step of production process is validated we can assure that the final product is of the best quality. Validation is best viewed as an important and integral part of GMP. Validation is therefore one element of quality assurance programs associated with a particular process. Quality cannot be assured only by doing finished product testing and in-process monitoring but it should be built into the manufacturing process. So building of quality require a special attention to a few factors like selection of material, process design, control variables, in process control and finished product testing. In this study three initial batch of clarithromycin tablet with same size, method, equipment and validation criteria were taken. The critical parameters involved in dry mixing, preparation of granulating agent, wet mixing, wet milling, drying, lubrication, and compression stages were identified and evaluated. Results obtained with this process validation data provides high degree of assurance that manufacturing process produce product meeting its predetermined specifications and quality attributes. It also provides documented evidence for the operation sequence of manufacturing process and to determine the critical parameters and variables in the process of manufacturing of the tablets. The output of process validation can be used to increase productivity, its consistent quality and decreasing the need for processing or market complaints.

**Keywords:** Clarithromycin, Process Validation, immediate release tablets, Control Variables.


User Name:

Password:

Remember Me

Search:



  
**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara





**BASIC ASPECTS OF PHARMACEUTICAL PROCESS VALIDATION OF SOLID DOSAGE FORMS: QUALITY ASSURANCE POINT OF VIEW**

**Ketaki S. Shinde<sup>1</sup>, Dr. R. C. Doijad<sup>1</sup>, Dr. J. S. Mulla<sup>1</sup> and Sachin S. Mali<sup>2</sup>**

<sup>1</sup>Department of Quality Assurance, Shree Santkrupa College of Pharmacy, Shivaji University, Ghogaon, India.

<sup>2</sup>Department of Pharmaceutics, Y. D. Mane Institute of Pharmacy, Kagal, India.

\*Corresponding Author: Ketaki S. Shinde

Department of Quality Assurance, Shree Santkrupa College of Pharmacy, Shivaji University, Ghogaon, India.

Article Received on 15/01/2022

Article Revised on 04/02/2022

Article Accepted on 25/02/2022

**ABSTRACT**

The purpose and interest of this overview on pharmaceutical process validation of immediate release tablets, is to highlight the critical process parameters to be validate during the activity of validation of solid dosage form. It is the most common dosage form for orally administration of drug. The Process validation should confirm that the control strategy is sufficient to support the process design and the quality of the product. This validation review covers the solid dosage form of process validation. The process is developed in such a way that the required parameters are achieved and it ensures that the output of the process will consistently meet the required parameters during routine production, the process is validated. A manufacturer can assure through careful design of the device, processes, process controls and process variable that all manufactured units will meet specifications and have uniform quality. This review provides information on objectives and benefits of process validation, types of process validation, major phases in validation and regulatory aspects. Guidelines and strategy for process validation of solid dosage form validation and regulatory aspects. Guidelines and strategy for process validation of solid dosage form are also discussed.

**KEYWORDS:** Pharmaceutical Process Validation, Process Validation Stages, Validation Acceptance Criteria.

**INTRODUCTION**

Validation is an integral part of quality assurance; it involves the systematic study of systems, facilities and processes aimed at determining whether they perform their intended functions adequately and consistently as specified. A validated process is one which has been demonstrated to provide a high degree of assurance that uniform batches will be produced that meet the required specifications and has therefore been formally approved. Validation in itself does not improve processes but confirms that the processes have been properly developed and are under control.<sup>[1]</sup> The objective of the design and manufacture of the immediate release tablet is to deliver orally the correct amount of drug in the proper form, over a period of time and in the desired location, and to have its chemical integrity protected to that point. Numerous features are required to ensure product quality and the validation is one of them. It is through careful design and validation of both the process and process controls that a manufacturer can establish a high degree of confidence that all manufactured units from successive lots will be acceptable. Successful validation of a process may reduce the dependence upon intensive in-process and finished product testing. In most cases, end-product testing plays a major role in assuring that quality assurance goals are meet. A validated process is one which has been demonstrated to provide a high

degree of assurance that uniform batches will be produced that meet the required specifications and has therefore been formally approved. Validation in itself does not improve processes but confirms that the processes have been properly developed and are under control.<sup>[2]</sup>

**Process Validation Definition<sup>[3]</sup>**

According to US FDA

In 1978

"A validation manufacturing process is one which has been proved to do what it purports or is represented to do. The proof of validation is obtained through the collection and evaluation of data, preferably, beginning from the process development phase and continuing the production phase. Validation necessarily includes process qualification (the qualification of materials, equipment, system, building, personnel), but it also includes the control on the entire process for repeated batches or runs"

In 1987

"Process validation is establishing documented evidence which provides a high degree of assurance that a specific process (such as the manufacture of pharmaceutical dosage forms) will consistently produce a product



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



ERWIN | ABSTRACT

ajmr.com/home/abstract.js?9508



**EUROPEAN JOURNAL OF  
PHARMACEUTICAL AND MEDICAL RESEARCH**  
(An ISO 9001:2015 Certified International Journal)

An International Peer Reviewed Journal for Pharmaceutical, Medical & Biological Sciences  
An Official Publication of Society for Advance Healthcare Research (Reg. No. - 01/0010161616)  
Medically Biotechnology, Global Impact Factor (GIF) (Under Process), Universal Impact Factor, Internal

ISSN 2094-3211  
IMPACT FACTOR: 6.222  
ICV - 79.57

SAFETY 2014 | 21:16:57 PM

ISO 9001:2015 Certified Journal

Login | Register

---

HOME | ABOUT US | INSTRUCTION TO AUTHOR | CURRENT ISSUE | MANUSCRIPT SUBMISSION | PROCESSING FEES | TRACK YOUR ARTICLE | ARCHIVE | CONTACT US

**Abstract**

**BASIC ASPECTS OF PHARMACEUTICAL PROCESS VALIDATION OF SOLID DOSAGE FORMS: QUALITY ASSURANCE POINT OF VIEW**

**Ketaki S. Shinde, Dr. R. C. Dojrad, Dr. J. S. Multa and Sachin S. Mali**

**ABSTRACT**

The purpose and interest of this overview on pharmaceutical process validation of immediate release tablets, is to highlight the critical process parameters to be validate during the activity of validation of solid dosage form. It is the most common dosage form for orally administration of drug. The Process validation should confirm that the control strategy is sufficient to support the process design and the quality of the product. This validation review covers the solid dosage form of process validation. The process is developed in such a way that the required parameters are achieved and it ensures that the output of the process will consistently meet the required parameters during routine production, the process is validated. A manufacturer can assure through careful design of the device, processes, process controls and process variable that all manufactured units will meet specifications and have uniform quality. This review provides information on objectives and benefits of process validation, scope of process validation, quality process is consistent and consistent quality.

SJIF Impact Factor: 6.222




**Citation:**

	All	Since 2018
Citation:	3096	2560

Type here to search

29°C Sunny 15:38 03-10-2024



  
**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## Review Article

**Floating Microspheres: A Novel Drug Delivery System**OMKAR B TIPUGADE <sup>1</sup>, JAMEEL AHMED S MULLA <sup>2\*</sup><sup>1</sup> Department of Pharmaceutics, Genesis Institute of Pharmacy, Radhanagari - 416212, Maharashtra, India<sup>2</sup> Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad - 415111, Maharashtra, India**ARTICLE DETAILS***Article history:*

Received on 8 November 2022

Modified on 6 December 2022

Accepted on 10 December 2022

*Keywords:*Floating Microspheres,  
Gastric Residence Time (GRT),  
Buoyancy.**ABSTRACT**

The motivation behind composing this audit on Floating microspheres is to gather the ongoing writing with unique spotlight on the essential system of buoyancy to accomplish gastric maintenance. Floating microsphere pledges to be a potential philosophy for gastric retention. The floating microspheres have been created trying to discharge the medication gradually into the GIT and keep up a compelling medication fixation in the serum for longer timeframe. From the system and technological factor of view, the floating drug delivery system is comparatively clean and logical approach. In this review, the current status of floating microspheres including hollow microspheres (micro balloons) and their characterization, advantages disadvantages, application, mechanism and method of preparation for gastric retention of drug are discussed. This review additionally summarizes the *in vitro* and *in vivo* studies to evaluate the overall performance and programs of floating microspheres.

© KESS All rights reserved

**INTRODUCTION**

The layout of oral managed drug delivery device have to in the main be geared toward achieving extra predictable and multiplied bioavailability of drug [1]. Thus the purpose of drug delivery gadget is to offer a therapeutic quantity of drug to the proper site within the frame to achieve promptly & then preserve the favored drug concentration. Drug which can be without difficulty absorbed from the GIT and having a brief half-life are eliminated fast from the blood circulation. To avoid these issues oral managed drug delivery structure were advanced as they releases the drug slowly into the GIT and maintain a regular drug concentration in the serum for longer duration of time [2]. It has been often determined that the drug that are easily absorbed form GI tract have a short half live and are eliminated quick from the systemic flow which result in incomplete of drug from upper a part of small intestine [3]. Recent clinical and patent literature has shown increased interest in novel dosage bureaucracy that can be retained in

the stomach for a extended and predictable duration of time. GRDF are designed on the idea of one of the several procedures like formulating low density dosage shape that stay buoyant above the gastric fluid (FDSD) or excessive density dosage shape that is retained at the bottom of the belly, imparting bio-adhesion to the stomach mucosa, reducing motility of the GIT by means of concomitant administration of medicine or pharmaceutical excipients, expanding the dosage shape via swelling or unfolding to a huge size which limits the emptying of the dosage form via the polymeric sphincter, making use of ion-alternate resin which adheres to mucosa, or using a modified shape system [4,5].

**Physiology Consideration:**

The belly is situated in the left upper part of abdominal cavity immediately under the diaphragm. Its size fluctuates in understanding to the measure of distension: as much as 1500 ml following a dinner; after nourishment has purged a fallen nation is gotten with resting amount of 25-50 ml. Anatomically the belly is split into three regions: fundus, frame and antrum (pylorus) [4,6]. The proximal part manufactured from fundus and frame acts as a reservoir for undigested material, the antrum is

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

181



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 14

October-December 2022

[← Back to Issue](#)

### Floating Microspheres: A Novel Drug Delivery System


Omkar B Tipugale, Jameel Ahmed S Mulla

[Download PDF](#)

**Abstract:**  
The motivation behind composing this audit on Floating microspheres is to gather the ongoing writing with unique spotlight on the essential system of buoyancy to accomplish gastric maintenance. Floating microsphere pledges to be a potential philosophy for gastric retention. The floating microspheres have been created trying to discharge the medication gradually into the GIT and keep up a compelling medication fixation in the serum for longer timeframe. From the system and technological factor of view, the floating drug delivery system is comparatively clean and logical approach. In this review, the current status of floating microspheres including hollow microspheres, matrix, balloon and their characterization, advantages

Type here to search 29°C Sunny 15:19 23-02-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara





## *Periplaneta americana* L. a potential source of traditional medicine: chemometric analysis, *in vitro* and *in silico* study

Suchismeeta Behera<sup>a,b</sup>, Amulyaratna Behera<sup>c</sup>, Suman Kumar Mekap<sup>c</sup>, Chinmaya Chidananda Behera<sup>c</sup>, Atul Kadam<sup>d</sup> and Prafulla K. Mohanty<sup>a</sup>

<sup>a</sup>Postgraduate Department of Zoology, Utkal University, Bhubaneswar, India; <sup>b</sup>State Forensic Science Laboratory, Bhubaneswar, India; <sup>c</sup>School of Pharmacy and Life Sciences, Centurion University of Technology and Management, Bhubaneswar, India; <sup>d</sup>Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, India

Communicated by Ramaswamy H. Sama

### ABSTRACT


Mayurbhanj is the ethnic dominant tribal population district in Odisha, India. The tribal's of Mayurbhanj depends on traditional medicines since time immemorial for health-related issues. Due to the imperative ethnic claim of traditional healers, the financial stringency of the patient community and the necessity to produce a better therapeutic effect has led to investigate ethno zoological sources and to find out the biochemical moiety responsible for the healing process. Considering the ethnic communities' acceptability of the zoological source as traditional medicine, the current evidence-based research study is conducted to investigate the biochemical moiety present in *Periplaneta americana*, responsible for therapeutic activity. The whole powdered *Periplaneta americana* was extracted using maceration techniques with n-hexane and methanol as solvent. The obtained extracts were subjected to GC-MS analysis to identify the biochemical moiety. To check the potential biological activity, an *in vitro* antimicrobial test was carried out in both turbidimetry and a viable count method against *E. coli*. Moreover, the obtained biochemical molecules were exposed to *in silico* studies for their binding modes and their affinity using Discovery studio software. The major compounds were found to be hexadecanoic acid, methyl ester, n-hexadecanoic acid, oleic acid, octadecanoic acid along with other minor constituents. The maximum inhibitory activity of n-hexane and methanol extract against *S. aureus* at a concentration of 400 µg/mL was found to be 89 and 87%, respectively. The binding models of almost all identified compounds confer very good binding affinities with some key and strong non-covalent interactions with various amino acid residues of receptor active site pocket, which predict the compounds to be potent inhibitors of various infectious bacteria. These findings suggested that the hexane extract of *P. americana* could be exploited as a potential natural source.

### ARTICLE HISTORY

Received 19 July 2020  
Accepted 27 May 2021

### KEYWORDS

*Periplaneta americana*;  
turbidimetry; viable count



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Periplaneta americana L. a pot... | sandfonline.com/doi/abs/10.1080/07391102.2021.1938681

Taylor & Francis Online | Log In | Register | Cart

Home | All journals | Journal of Biomolecular Structure and Dynamics | List of Issues | Volume 40, Issue 20 | Periplaneta americana L. a potential sou...

Journal of Biomolecular Structure and Dynamics > Volume 40, 2022 - Issue 20

Submit an article | Journal homepage

Enter keywords, authors, DOI, etc. | The Journal | Advanced search

129 Views  
0 CrossRef citations to date  
0 Attempts

Research Articles

### Periplaneta americana L. a potential source of traditional medicine: chemometric analysis, *in vitro* and *in silico* study

Suchismeeta Behera, Amulyaratna Behera, Suman Kumar Meikap, Chinmaya Chidananda Behera, Abul Kadam & Prafulla K. Mohanty


Pages 1931-1947 | Received 19 Jul 2020; Accepted 27 May 2021; Published online 21 Jun 2021

Cite this article | <https://doi.org/10.1080/07391102.2021.1938681>

Full Article | Figures & data | References | Citations | Metrics | Reprints & Permissions | Read the article

Type here to search | 29°C Sunny | 15:23 03-02-2024



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



## Review Article

**Non-Ionic Surfactant Vesicle (Niosome): A Novel Drug Delivery System**

SANDHYA S SHEWALE, JAMEEL AHMED S MULLA\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Dist: Satara, MS - 415111, India

**ARTICLE DETAILS***Article history:*

Received on 6 August 2022

Modified on 24 September 2022

Accepted on 27 September 2022

*Keywords:*

Non Ionic Surfactant Vesicle,

Niosomes,

Preparation,

Characterization,

Applications.

**ABSTRACT**

Niosomes are vesicles that are created by hydrating a mixture of lipids that are biodegradable, non-ionic surfactant, and cholesterol. In comparison to a drug's traditional dosing form, niosomes boost the drug's action. Drugs that are amphiphilic or lipophilic can be transported via niosomes. The problems associated with pharmaceutical instability, rapid disintegration, insolubility, and low bioavailability may be resolved by niosomes. The manner of formulation determines whether niosomes are multilamellar or unilamellar in structure. For the site-specific administration of anti-cancer, anti-infective drugs, etc., niosomes have a very effective drug delivery capability. In comparison to other drug formulations, niosomes are stable and inexpensive carriers. Niosomes are also used in innovative drug delivery systems, topical drug delivery systems, oral drug delivery systems, and parental drug delivery systems. This review provides an extensive summary of niosomal studies to date, as well as a detailed look at formulation aspects, niosome types, physical characterization methods, and recent pharmaceutical applications like transmucosal, oral, ocular, topical, and pulmonary drug delivery as well as cosmetic applications.

© KESS All rights reserved

**INTRODUCTION**

"A good drug delivery system delivers drug to the T-site for use throughout treatment. Targeted delivery is a strategy for delivering drugs to tissues while reducing the concentration of the drug in surrounding tissue. Niosomes are thin lamellar vesicles that are not toxic to organisms. They are formed by mixing cholesterol with a non-toxic alkyl class surfactant or dialkylpolyglycerol ether, followed by hydration in water [1]. Niosomes are versatile and can be combined to deliver several forms of drugs to the intended site. A niosome is a vesicle that can be unilamellar or multilamellar and is formed of non-ionic surfactant, cholesterol, and ionic surfactant. Its purpose is to reduce structural bonds. Drugs that are hydrophilic, lipophilic, or amphiphilic can be incorporated into the niosome's bilayer structural vesicle. Niosome shows more stability than liposome because the liposome can be degraded and oxidized due to its particular lipophilic nature.

Because niosomal formulations have a non-ionic surfactant, they remain in the bloodstream for a longer period of time, enhancing their target action [2,3].

**Structure of Niosomes**


Niosomes are microscopic, spherical, lamellar (unilamellar or multilamellar) structures. The bilayer is produced by combining charge-inducing agent with nonionic surfactants, either with or without cholesterol. Niosomes are formed by combining various surfactant types in different combinations and molar ratios. Alkyl glyceryl ethers, Alkyl ethers, polyoxyethylene fatty acid esters and sorbitan fatty acid esters are a few examples of surfactants [2,4]. Cholesterol addition keeps the bilayer firm, resulting in fewer leaky niosomes [5]. Charge inducers, but on the other hand, give the vesicles a charge and expand their size, improving drug entrapment effectiveness. The vesicles are stabilised by positive charge inducers such stearylamine and cetylpyridinium chloride as well as negative charge inducers like dihexadecyl phosphate, dicetyl phosphate, and lipoamino acid [6]. Nonionic surfactants in niosomes have a tendency to have hydrophobic ends that face

**\*Author for Correspondence:**

Email: jameelahmed5@rediffmail.com

129



  
Principal  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

Non-ionic surfactant vesicle (niosome): A novel drug delivery system

Indian Journal of Novel Drug Delivery

ISSN 0975-5500

IDAAM Publications

Home About Articles & Issues Publish Subscriptions Advertise Contact

VOLUME 14

July-September 2022


Non-ionic surfactant vesicle (niosome): A novel drug delivery system

Sandhya S Shewale, Jameel Ahmed S Mulla

Download PDF

**Abstract:**  
Niosomes are vesicles that are created by hydrating a mixture of lipids that are biodegradable, non-ionic surfactant, and cholesterol. In comparison to a drug's traditional dosing form, niosomes boost the drug's action. Drugs that are amphiphilic or lipophilic can be transported via niosomes. The problems associated with pharmaceutical instability, rapid disintegration, insolubility, and low bioavailability may be resolved by niosomes. The manner of formulation determines whether niosomes are multilamellar or unilamellar in structure. For the site-specific administration of anti-cancer, anti-infective drugs, etc. niosomes have a very effective drug delivery capability in comparison to other drug formulations. niosomes are stable and



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara





**REVIEW ARTICLE**

**Nanosuspension: A Novel Technology for Drug Delivery**

Vidya Ashok Kheradkar, Jameel Ahmed S. Mulla\*

Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, MS-415111, India.

\*Corresponding Author E-mail: [jameelahmed5@rediffmail.com](mailto:jameelahmed5@rediffmail.com)

**ABSTRACT:**

Nanotechnology includes nanosuspensions. Nanotechnology has become a huge area in the world of medicine. An appealing and promising approach to address the lipophilic like bio-available and poor solvability concerning the pharmaceuticals is nanosuspension. The majority of recently developed medications have poor water solubility. Drug formulation is usually complicated by medicines' limited water solubility. There are numerous standard methods for improving the solubility of pharmaceuticals that are poorly soluble, including Media Milling, Homogenization, precipitation technique, et cetera. A new approach in medication development is the development disintegrable medicinal formulations using nanotechnology. Particle sizes used in nanotechnology range are from 1 to 1000 nm. The increase in dissolving rate that results from drug particle size reduction into the submicron region naturally improves bioavailability. An efficient new method for improving the dissolution of low soluble in water medicines is nanosuspension technology. A nanosuspension improves the safety and effectiveness of a medicine by addressing issues with lipophilic like bio-available and poor solvability, as well as by changing the pharmacokinetics.

**KEYWORDS:** Nanosuspension, Preparation methods, Characterization, Evaluation, Applications.

**INTRODUCTION:**

All branches of applied science, manufacturing, and production are embracing nanotechnology. It is a brand-new field of thorough intensive studies that integrates medicine and other life sciences. The high technology and advanced science of nanotechnology precisely defines the molecular structure of matter<sup>1</sup>. Colloidal dispersions and biphasic systems called nanosuspensions are composed of particles distributed in an aqueous medium and whose diameter is smaller than 1  $\mu$ m. In nanosuspensions solidified atom typically have a particle size distribution that is smaller than one micron<sup>1,2</sup>.

It allows the prospect for fresh and distinctive techniques with several different applications in the treatment of cancer, including diagnostics, therapy, and prognosis. The enhanced chemical characteristics and physical of nanoscale particles are their principal benefit.


Particle size, surface area, hydrophobicity, crystallinity, and surface charge are some of the key factors in drug delivery<sup>2,3</sup>.

Nanoparticles and solid nanoparticles of lipid are two different types of nanosuspensions. Solid nanoparticles in lipid are lipid carriers of pharmaceuticals, whereas nanoparticles are often polymeric colloidal transporters of drugs<sup>3,4</sup>. These can be utilised to increase the soluble amount of medications that are insufficiently soluble in lipid and aqueous media. The biopharmaceutical categorization system (BCS) classes II and IV are all drug molecules that can use the nanosuspension formulation to make them more soluble<sup>4</sup>.

To increase the solubility of pharmaceuticals that are poorly soluble in water, numerous conventional methods have been developed, including micronization, solubilization employing co-solvents, surfactant dispersions, and precipitation technique. Intravenous fluid, gastrointestinal, ophthalmic, and respiratory routes can all be used to administer nanosuspensions<sup>4,5</sup>. At the moment, work is focused on expanding their uses for site-specific medication delivery<sup>5</sup>.

Received on 14.10.2022 Modified on 16.01.2023  
Accepted on 12.04.2023 © Asian Pharma Press All Right Reserved  
Asian J. Res. Pharm. Sci. 2023; 13(2): 106-110.  
DOI: 10.52711/2231-5659.2023.00020



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



Asian Journal of Research in Pharmaceutical Sciences

ISSN 2231-5659 (Online)  
2231-5640 (Print)

ABOUT JOURNAL EDITORIAL BOARD FOR AUTHORS MORE NEWS

SEARCH Submit Article

## Nanosuspension: A Novel Technology for Drug Delivery

Author(s): Vidya Ashok Kheradkar, Jameel Ahmed S. Mulla  
 E-mail(s): jameelahmed5@gmail.com  
 DOI: 10.32711/2231-5659/2023.00020

Address: Vidya Ashok Kheradkar, Jameel Ahmed S. Mulla\*  
 Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, MS-415131, India.  
 \*Corresponding Author

Published in: Volume - 13, Issue - 2, Year - 2023

Purchase PDF View HTML

Asian Journal of Research in Pharmaceutical Sciences (AJPSci.) is an international, peer-reviewed journal, devoted to pharmaceutical sciences.

Read more

PDF Not Available  
 DOI: 10.32711/2231-5659



**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## Original article

## Evaluation of solid-lipid nanoparticles formulation of methotrexate for anti-psoriatic activity



Debarati Maiti<sup>a</sup>, Mohammed Naseeruddin Inamdar<sup>a,b</sup>, Mansour Almuqbil<sup>c</sup>, Sarasija Suresh<sup>d</sup>, Syed Mohammed Basheeruddin Asdaq<sup>e,\*</sup>, Sultan Alshehri<sup>f</sup>, Saad Ali Al Arfaj<sup>g</sup>, Ali Musharraf Alamri<sup>h</sup>, Meshal Meshary Aldohyan<sup>i</sup>, Misfir Theeb Alqahtani<sup>j</sup>, Turki Mohammed Alosaimi<sup>k</sup>, Sami Haran Alenazi<sup>l</sup>, Moneer E. Almadani<sup>m</sup>, Jameel Ahmed S. Mulla<sup>n</sup>, Syed Imam Rabbani<sup>o</sup>

<sup>a</sup> Department of Pharmacology, Al-Ameen College of Pharmacy, Bangalore, India

<sup>b</sup> Department of Pharmacology, East West College of Pharmacy, Bangalore, India

<sup>c</sup> Department of Clinical Pharmacy, College of Pharmacy, King Saud University, Riyadh 11451, Saudi Arabia

<sup>d</sup> RGV Research and Innovations Pvt Ltd (RGIIR), Bangalore, India

<sup>e</sup> Department of Pharmacy Practice, College of Pharmacy, AlMaarefa University, Daryyah, 13713, Riyadh, Saudi Arabia

<sup>f</sup> Department of Pharmaceutical Sciences, College of Pharmacy, AlMaarefa University, Ad Diryah 13713, Saudi Arabia

<sup>g</sup> King Abdulaziz Medical City in Riyadh, Ministry of National Guard, Riyadh, Saudi Arabia

<sup>h</sup> King Salman Hospital, Riyadh, Saudi Arabia

<sup>i</sup> Department of clinical medicine, College of medicine, AlMaarefa University, Daryyah, Riyadh 13713, Saudi Arabia

<sup>j</sup> Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Maharashtra, India

<sup>k</sup> Department of Pharmacology and Toxicology, College of Pharmacy, Qassim University, Buraydah 51452, Saudi Arabia

## ARTICLE INFO

Article history:  
Received 5 March 2023  
Accepted 6 April 2023  
Available online 15 April 2023

Keywords:  
Solid-lipid nanoparticles  
Methotrexate  
Permeability  
Distribution  
Keratinocytes

## ABSTRACT

**Background & Objectives:** Methotrexate (MTX) is commonly used to manage psoriasis. The drug has erratic absorption characteristics and shows several complications. The present study uses different experimental models to evaluate the solid-lipid nanoparticles of MTX (SLN-MTX) for the anti-psoriatic effect. **Methods:** A prepared SLN-MTX formulation was used and its permeability studies were conducted on Wistar rat abdominal skin. The organ-level distribution of the drug in the formulation was tested in mice and the *in-vitro* anti-psoriatic activity was determined in CL-177; XB-2 keratinocytes cell lines. The efficacy of SLN-MTX formulation was compared with standard MTX and marketed MTX preparations. The results are analyzed statistically using the student's *t*-test.

**Results:** The data suggested that MTX from the formulation was slowly released and completely (80.36%) permeated through the skin. The flux and permeation data were found to be maximum for SLN-MTX compared to marketed and standard preparations. MTX in the formulation was found to be distributed more in the liver (67.5%) and kidney (2.34%). Further, SLN-MTX formulation showed dose-dependent inhibition on the growth of keratinocytes, and the cytotoxic concentration (CTC50) was found to be the least (518 mcg/ml).

**Interpretation & Conclusion:** The findings suggested that MTX in solid-lipid nanoparticles could be a promising formulation for the management of psoriasis since the drug was slowly released, progressively

\* Corresponding author at: Department of Pharmacy Practice, College of Pharmacy, AlMaarefa University, Daryyah, 13713, Riyadh, Saudi Arabia.

E-mail addresses: ojalthyasadq@gmail.com (D. Maiti), inamdam@gmail.com (M. Naseeruddin Inamdar), moneerwari@ksu.edu.sa (M. Almuqbil), sarasija\_s@hotmail.com (S. Suresh), asdaq9@gmail.com, sasdaq@mcst.edu.sa (S. Mohammed Basheeruddin Asdaq), alshehri@mcst.edu.sa (S. Alshehri), arfaj@nguhj.med.sa (S. Ali Al Arfaj), meshalme@nguhj.med.sa (A. Musharraf Alamri), imnd444@hotmail.com (M. Meshary Aldohyan), misfir\_gh@hotmail.com (M. Theeb Alqahtani), researcher1822@gmail.com (T. Mohammed Alosaimi), Sami-5@hotmail.com (S. Haran Alenazi), researcher072022@gmail.com, researcher172022@gmail.com (M.E. Almadani), jameelahmed5@rediffmail.com (J. Ahmed S. Mulla), syedimamrabbani@yahoo.com (S. Imam Rabbani).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

<https://doi.org/10.1016/j.sps.2023.04.007>

1319-0164/© 2023 The Author(s). Published by Elsevier B.V. on behalf of King Saud University.  
This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara

ScienceDirect Journals & Books Search... Register Sign In

View PDF Download full issue

Outline  
 Abstract  
 Keywords  
 1. Introduction  
 2. Methods  
 3. Results  
 4. Discussion  
 5. Conclusion  
 Declaration of Competing Interest  
 Acknowledgment  
 Funding  
 References  
 Show full outline

Saudi Pharmaceutical Journal  
 Volume 32, Issue 6, June 2021, Pages 834–844

Original article

### Evaluation of solid-lipid nanoparticles formulation of methotrexate for anti-psoriatic activity

Debarati Maji<sup>a</sup>, Mohammed Nawaaruddin Inamdar<sup>b</sup>, Mansour Almuqbil<sup>c</sup>,  
 Somnath Suresh<sup>d</sup>, Saeed Mohammed Basheeruddin Adooq<sup>e</sup>, Sultan Alshelhi<sup>f</sup>,  
 Saad Ali Al Arfaj<sup>g</sup>, Ali Mubhammad Alami<sup>h</sup>, Meshal Meqhar Aladwan<sup>i</sup>,  
 Wasif Thash Alaghtani<sup>j</sup>, Taki Mohammed Alqasbi<sup>k</sup>, Sami Hassan Alengal<sup>l</sup>,  
 Moneer E. Almadani<sup>m</sup>, Jameel Ahmed S. Mulla<sup>n</sup>, Saeed Imran Rabbani<sup>o</sup>

Recommended articles  
 Community pharmacist's preparedness to provide patient-centred care in Saudi...  
 Saudi Pharmaceutical Journal, Volume 31, Issue 4, ...  
 Muhammad Karim Basheer, ... Zabeer Us-Din Boker  
 View PDF  
 Design, physicochemical characterisation, and in vitro cytotoxicity...  
 Saudi Pharmaceutical Journal, Volume 31, Issue 6, ...  
 Muhammad H. Sultan, ... Syam Mohan  
 View PDF  
 Benzofurquinazolines as antifungal against candidiasis: Screening, molecu...  
 Saudi Pharmaceutical Journal, Volume 31, Issue 6, ...  
 Husein A. Abuoliz, ... Rashid A.  
 FEEDBACK



**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



**Cubosomal Drug Delivery System: Promising Approach for Ocular Drug Delivery of Corticosteroid**  
**Snehal Chakorkar<sup>b</sup>, Jameel Ahmed S. Mulla<sup>a</sup>**

<sup>a</sup> Department of Pharmaceutics, Shree Santkrupa College of Pharmacy, Ghogaon, Karad

<sup>b</sup> Department of Pharmacology, Dr. D. Y. Patil Institute of Pharmaceutical Sciences and Research, Pimpri, Pune.

\*Corresponding Author: Dr. Jameel Ahmed S. Mulla

[jameelahmed5@gmail.com](mailto:jameelahmed5@gmail.com), Professor & Head, Department of Pharmaceutics

Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Satara

Maharashtra – 415111 India

**Abstract:**

**Background:** Over a century ago, uveitis, sclerokeratitis, pemphigus, and keratoconjunctivitis were treated with systemic administration of hypophyseal-derived adrenocorticotrophic hormone. Although expanding research has effectively improved the use of corticosteroids locally, various side effects prevent patient compliance. Scientists are continuously developing different strategies and logical and dynamic approaches to enhance the acceptance of corticosteroid drug delivery.

**Objective:** Over recent decades, the number of problems associated with using corticosteroids was overcome by changing the route of systemic drug delivery to ocular drug delivery. The recent development of lipid-based drug delivery has made it possible to enhance the bioavailability of corticosteroid ocular drug delivery. These novel methods may overcome the ocular barriers and side effects of conventional eye drops. Many of these formulations are presently undergoing clinical trials, and few are clinically effective.


**Result:** This review presents an overview of corticosteroid-loaded cubosomal drug delivery as a non-invasive, nanoparticulate topical ocular delivery system.

**Conclusion:** This novel formulation is easy to formulate, has a high residence time, sustains drug release, and enhances the bioavailability of corticosteroids. On this basis sheds light on the anatomy of the eye, various barriers to ocular drug delivery, various approaches for ocular drug delivery, mechanism of cubosomal absorption, polymers used,

13850

Eur. Chem. Bull. 2023, 12(Special Issue 4), 13850-13881



  
**Principal**  
**Dr. Ramling G. Patrakar**  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



European Chemical Bulletin

www.chemicalbulletin.com/issue-content/subsonal-drug-delivery-system-promising-approach-for-ocular-drug-delivery-of-corticosteroid-5440

ISSN 2093-5340

Search for title, author, etc

Submit article

European Chemical Bulletin

ISSN 2093-5340

Open Access Journal

BIOCHEMISTRY OF FASTING – A REVIEW ON METABOLIC SWITCH AND AUTOPHAGY.

ONE-JOY ENVIRONMENT 123456789

Cubosomal Drug Delivery System: Promising Approach for Ocular Drug Delivery of Corticosteroid

PDF

Keywords: Cubosomes, Ocular drug delivery, Corticosteroid, Novel drug delivery

Snehal Chakrkar, Jameel Ahmed S. Mulla  
+ doi: 10.48847/eeb-2023-12-s4-1258

Abstract

Over a century ago, uveitis, sclerokeratitis, pemphigus, and keratoconjunctivitis were treated with systemic administration of hypophyseal-derived adrenocorticotropic hormone. Although expanding research has effectively improved the use of corticosteroids locally, various side effects prevent patient compliance. Scientists are continuously developing different strategies and logical and dynamic approaches to enhance the acceptance of corticosteroid drug delivery. Objective: Over recent decades, the number of problems associated with using corticosteroids was overcome by changing the route of systemic drug delivery to ocular drug delivery. The recent

ELSEVIER  
ORCID  
Crossref  
OPEN ACCESS

Type here to search

39°C Sunny

15:31  
23-10-2024



**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



## Phytochemical and Pharmacological Review on *Garcinia talbotii* : A Latent Medicinal Plant

Abhishek Vilas Desai<sup>1</sup> and Jameel Ahmed S. Mulla<sup>2\*</sup>

<sup>1</sup>Research Scholar, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Maharashtra, India and

<sup>2</sup>Assistant Professor, Ashokrao Mane Institute of Pharmacy, Ambap, Kolhapur, Maharashtra, India

<sup>3</sup>Professor, Shree Santkrupa College of Pharmacy, Ghogaon, Karad, Maharashtra, India.

Received: 01 July 2023

Revised: 20 Aug 2023

Accepted: 21 Sep 2023

### \*Address for Correspondence

Jameel Ahmed S. Mulla

Professor,

Shree Santkrupa College of Pharmacy,

Ghogaon, Karad, Maharashtra, India.

E.Mail: jameelahmed5@rediffmail.com



This is an Open Access Journal / article distributed under the terms of the **Creative Commons Attribution License (CC BY-NC-ND 3.0)** which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. All rights reserved.

### ABSTRACT

*Garcinia talbotii* Raizada ex Santapau commonly known as "Tavir" in India, is an evergreen dioecious medium sized tree distributed in the semi evergreen to evergreen forests of Western Ghats. Leaves elliptic-oblong or ovate, 5–18 × 4–9 cm. Flowers creamy-white, white to greenish-yellow Fruits globular, 4–6 cm. Traditionally Dried fruits are used like tamarind in curries. *Garcinia* species has a lot of medicinal properties but Limited research work has been carried out for *G. talbotii* and scanty reports are available about phytochemical investigation and assessment of different biological activities, so a review work was carried out to study its phytochemical properties and biological activities.


**Keywords:** *Garcinia talbotii*, Phytochemical, Phytoconstituents, Characterization, Pharmacological screening


### INTRODUCTION


Due to the presence of two key bioactive constituents, hydroxycitric acid and garcinol, *Garcinia* L. (Clusiaceae) is one of the medicinally and commercially noteworthy genera. There are 260 species in the genus, with the majority of them found in tropical areas, notably Asia [1]. With some species ranging as far as India and the Micronesian islands, as well as tropical Africa and the Neotropics, the Malaysian region is thought to be the centre of *Garcinia* diversity.[2] *Garcinia* has 39 species and 7 variants, of which 18 species and 5 varieties are native to India.[3] *Garcinia* species have received considerable attention worldwide from the scientific as well as industrial sectors due to the report of several bioactive structures such as biflavonoids, xanthenes and benzophenones.[4] Economically, the genus is crucial to several businesses, including food, modern medicine, and agriculture. The genus is significant because it

62391



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



  
**Principal**  
**Dr. Ramling G. Patrakar**  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara



# Formulation and Evaluation of Poly Herbal Chewable Tablets from Ayurvedic Ingredients having Cough Relieving Activity

Vinayak Dasharath Gaikwad\*  
Shri. Santkrupa College of Pharmacy,  
Ghogaon, Karad, Satara,  
Shivaji University, Kolhapur,  
Maharashtra 415111, India.

Priyanka A. Thorat (Assistant Professor)  
Department of Pharmaceutics,  
Shree Santkrupa College of Pharmacy,  
Ghogaon, Satara, Shivaji University, Kolhapur,  
Maharashtra 415111, India

**Abstract:-** In recent years, there is a spurt in the interest regarding use of Ayurvedic forms of medication. Glycyrrhiza Glabra (Licorice), Zingiber Officinale (Ginger), Curcuma longa (Turmeric) has been used as medicine from ancient times for treating the cough. These herbal drugs use was mentioned in Ayurveda. Cough affects patients of every age, so it is the common disease condition. Because of better patient compliance, flexible design of dosage forms, ease of administration and least sterility requirements, oral route of drug administration is the well liked route. The goal of this research work is to develop Polyherbal chewable tablets of various Ayurvedic Drugs by wet granulation method and to evaluate the formulations for various pharmaceutical parameters. The purpose of this study was to produce polyherbal chewable tablets using Licorice, Ginger, Turmeric. These polyherbal chewable tablets were formulated by technique of wet granulation method by utilizing binder that is acacia gum (5% w/v). Quality of final polyherbal chewable tablet was evaluated for pre formulation and post formulation parameter. Bulk and tapped densities, angle of repose, Carr's index, and Hausner's ratio are the pre-formulation tests evaluated for the prepared powder mixture (blend). Polyherbal chewable tablets were examined for general appearance, tablet size and shape, hardness, friability, weight variation as well as duration of disintegration.

**Keywords:-** Polyherbal Chewable Tablet; Glycyrrhiza Glabr; Licorice; Zingiber Officinal; Ginger; Curcuma longa; Turmeric; Wet granulation method; Cough; Antitussive; Demulcent; Expectorant; Cough relieving activity; Disintegration Test.

## I. INTRODUCTION

Several pharmacological cough relievers are sold in the medical stores and generally taken for the cough treatment. Now a days, the herbal cough relieving agents use are increasing day by day due to the high quality effectiveness with less side effects. Chewable tablets are a convenient dose form that have multiple good factors that is the strength and firmness of solid medications with the convenience of consuming and medication administration with no necessity for water.

Ayurvedic medicines have been used for several centuries for both the avoidance and cure of coughing. Although many over-the-counter medications are accessible in all medical stores, neither antiviral nor antibiotic therapy has been demonstrated to be treatable in the absence of initial lung infection.


These formulated polyherbal chewable tablets contains herbal Crude drugs such as Licorice (Glycyrrhiza Glabra), Turmeric (Curcuma longa) and Ginger (Zingiber Officinale). Chewable tablets are available in several different kinds of sizes and shapes. Chewable tablets are an oral delivery system commonly utilised for pharmaceutical and nutraceutical products. Chewable tablets that are chewed can have complete weights that are higher than those formulated as ordinary swallowing tablets, that are typically lower weight than 1g. Before swallowing, chewable tablets are crushed in the oral cavity. Biting of the chewable tablets starts in the mouth, the tablet is squashed or crushed into mouth and converted to the littler particles from where increment dissolution and ensuing better availability occurs to give the ideal therapeutic or pharmacological activity.

Chewable tablets is an fast releasing medication that are taken orally and chewed by the patient's teeth before being engulfed completely.[1]

The chewable tablets have an even, flat surface and a mild flavor without harsh and terrible taste; they are recommended to be disintegrating in the oral cavity more gradually, whether chewing is occurring or not. These chewable tablets not contains any flavouring agent and sweetener. The successful formulation development depends on the appropriate excipients selection. Chewable tablets are of the large size which are hard for gulping accordingly, chewable tablets bit in buccal cavity before swallowing.[2]

The expanded bioavailability of chewable tablets occurs because of expanded absorption characteristics. Its disintegration or being bitten in the mouth forms littler molecules, for example, expanded surface zone of drug molecules of tablet into Gastrointestinal tract. Chewable Tablets have this advantage over ordinary capsules or strong (swallowing) tablets which are ingested predominantly after dissolution and breaking down. Chewable tablets are defined as palatable medicines that can be broken down and



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



INTERNATIONAL JOURNAL OF INNOVATIVE SCIENCE AND RESEARCH TECHNOLOGY

**IJISRT A DIGITAL LIBRARY**

ISSN NO : 2456-2165

ABOUT US | CALL FOR PAPERS | TOPICS | FOR AUTHORS | SPECIAL ISSUE | BROWSE ARCHIVE | PEER REVIEWS | CONTACT US | SIGN UP | LOGIN

Enter Name of Paper

Formulation and Evaluation of Poly Herbal Chewable Tablets from Ayurvedic Ingredients having Cough Relieving Activity

Download PDF Full text version

Apply For Certificate Hard Copy

Authors : Vinayak Dadasarath Gokwad, Priyanka A. Thorat

Volume/Issue : Volume 8 - 2023, Issue 4 - April

Google Scholar : <https://bit.ly/3TmGsDi>

CALL FOR PAPERS

Paper Submission Last Date  
29 - February - 2024

Paper Review Notification  
In 1-2 Days


Paper Publishing  
In 2-3 Days

SUBMIT YOUR PAPER

Type here to search

30°C Sunny 15:59 23-02-2024



  
**Principal**  
 Dr. Ramling G. Patrakar  
 Shree Santkrupa College of Pharmacy  
 Ghogaon, Tal. Karad, Dist. Satara

RESEARCH ARTICLE

## Extraction, Phytochemical Screening and HPTLC Analysis of Fenugreek Gum for Analysis of Different Phytoconstituents.

Pravinkumar D Lade<sup>1,2\*</sup>, Neelam Singla<sup>1</sup>

<sup>1</sup>Suresh Gyan Vihar University Jaipur Rajasthan, India

<sup>2</sup>Shree Santkrupa College of Pharmacy, Satara, Maharashtra, India.

Received: 10<sup>th</sup> October, 2023; Revised: 12<sup>th</sup> November, 2023; Accepted: 07<sup>th</sup> December, 2023; Available Online: 25<sup>th</sup> December, 2023

### ABSTRACT

Fenugreek gum is a natural polysaccharide that has been explored for various biomedical applications. The present study investigated the phytoconstituent profile of fenugreek gum, revealing it as a potent source of diverse bioactive compounds. The analysis encompassed steroids, fats and oils, alkaloids, proteins, saponins, tannins, phenolics, flavonoids, and carbohydrates. Extraction using methanol and aqueous solvents demonstrated the highest concentration of phytoconstituents, underscoring the efficacy of these solvents in isolating bioactive components from fenugreek. Further exploration involved the determination of Rf values and percentage area in chloroform, petroleum ether, and chloroform extracts, providing a comprehensive identification of the mixture's components. The results suggested a rich and varied chemical composition in fenugreek gum, offering potential applications in diverse fields owing to the presence of steroids for physiological effects, fats and oils for nutritional benefits, alkaloids with potential pharmacological properties, proteins for cellular functions, saponins for their soap-like properties, and tannins and phenolics as antioxidants. Additionally, the presence of flavonoids and carbohydrates adds to its nutritional and medicinal value. The study's findings contribute valuable insights into the phytochemical composition of fenugreek gum, shedding light on its potential applications in the realms of medicine, nutrition, and industry. The identification of optimal extraction methods and the characterization of phytoconstituents lay the foundation for future research and utilization of fenugreek gum as a versatile and valuable natural resource. Overall, this investigation underscores the significance of fenugreek gum as a rich repository of bioactive compounds with diverse potential applications across various domains.

**Keywords:** Fenugreek gum, HPTLC, Phytochemicals, Extraction, Herbal, Natural gum.

International Journal of Pharmaceutical Quality Assurance (2023), DOI: 10.25258/ijpqa.14.4.42

**How to cite this article:** Lade PD, Singla N. Extraction, Phytochemical Screening and HPTLC Analysis of Fenugreek Gum for Analysis of Different Phytoconstituents. International Journal of Pharmaceutical Quality Assurance. 2023;14(4):1090-1093.

**Source of support:** Nil.

**Conflict of interest:** None

### INTRODUCTION

Fenugreek (*Trigonella foenum graecum*) is an annual plant classified under the Leguminosae or Fabaceae family. Its seeds are composed of a high amount of soluble fibers which is also known as galactomannans. This plant is very potential in terms of herbal remedy due to the presence of wider phytoconstituents.<sup>1</sup> The gums derived from this plant has potential benefits in the pharmaceutical industry. Galactomannans obtained from different sources have different physico chemical properties. They may have different molecular weights and solubility profiles.<sup>2</sup> The distinctive feature of fenugreek gum lies in its equal galactose:mannose ratio of 1:1, showcasing the highest galactose concentration. This property results in increased solubility in water as compared to guar and locust bean gum. The greatest hydration and solubility of fenugreek gum are attributed to the homogeneous linking of its galactose and mannose moieties. Galactomannan physicochemical qualities are significantly influenced by the M/G ratio, with


solubility exhibiting an inverse relationship with this ratio. This characteristic leads to enhanced aqueous solubility in comparison to the other gums. The uniform linkage of galactose and mannose in this gum is responsible for its higher solubilities and hydration properties. The M/G ratio influences the physicochemical properties of galactomannans, with solubility showing an inverse correlation to this ratio.<sup>3</sup>


The M/G ratio plays a pivotal role in influencing the physicochemical properties of galactomannans, exhibiting an inverse relationship with the gum's solubility.<sup>4</sup> By adhering to oil droplets, galactomannans help to stabilize the steric balance of emulsions by inhibiting flocculation and coalescence. These biopolymers also exhibit emulsification, interfacial, and surface activity. Fenugreek gum is favored above other natural hydrocolloids because of these unique qualities, which make it a great component for many culinary applications.<sup>5</sup>


Fenugreek gum imparts desirable characteristics such as texture, appeal, gelling, thickening, and emulsifying properties.

\*Author for Correspondence: pldode78@gmail.com



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara



  
**Principal**  
Dr. Ramling G. Patrakar  
Shree Santkrupa College of Pharmacy  
Ghogaon, Tal. Karad, Dist. Satara